

SECTION : (circle one)

NAME : _____
Arkowitz Vatter

Math 8

March 16, 2009

Final Exam

INSTRUCTIONS: This is a closed book exam and no notes are allowed. You are not to provide or receive help from any outside source during the exam except that you may ask the instructor for clarification of a problem. You have three hours and you should attempt all problems.

- *Print* your name in the space provided and circle your instructor's name.
 - Calculators or other computing devices are not allowed.
 - Use the blank page at the end of the exam for scratch work.
 - **Except in the multiple choice section, you must show all work and give a reason (or reasons) for your answer. A CORRECT ANSWER WITH INCORRECT WORK WILL BE CONSIDERED WRONG.**
-
-

1. (10) Express the integral

$$\int \frac{\sin x - x}{x} dx$$

as an infinite series. Your answer should be in the form $\sum_{n=0}^{\infty} c_n x^n$, for some coefficients c_n .
(In other words, do *not* simply give the first few terms.)

2. (16) Determine the integral

$$\int \frac{\sqrt{x^2 - 1}}{x^3} dx.$$

3. (14) Determine the integral

$$\int \frac{x^2}{2} f''(x) dx.$$

Your answer should contain $f(x)$, $f'(x)$ and $\int f(x) dx$.

4. (16) A particle moves in space and its acceleration at time t is given by

$$a(t) = \langle e^t, 0, e^{-t} \rangle.$$

Its initial velocity is $v(0) = \langle 1, \sqrt{2}, -1 \rangle$ and its initial position is $s(0) = \langle 3, 0, 2 \rangle$.

(a) What is the particle's velocity and position at any time t ?

(b) What is the distance the particle has moved (along its path) from time $t = 0$ to time $t = \ln 2$. Simplify your answer by expressing it as a decimal.

5. (10) Let $w = xy + z$ and let $x = s \cos t$, $y = \ln s$ and $z = \arctan(s + t)$. Find

$$\frac{\partial w}{\partial s}.$$

6. (16) The function $z = f(x, y)$ has, at the point $(1, 2)$, $D_{\langle 1/\sqrt{2}, 1/\sqrt{2} \rangle}(1, 2) = 7/\sqrt{2}$ and $D_{\langle 1/\sqrt{5}, 2/\sqrt{5} \rangle}(1, 2) = 11/\sqrt{5}$. What is the directional derivative at $(1, 2)$ in the direction from $(1, 2)$ to $(3, 5)$?

7. (14) Find the equation of the tangent plane to the surface defined by

$$x - z = 4 \arctan(yz)$$

at $(1 + \pi, 1, 1)$.

8. (14) Determine the local maximum point(s), local minimum point(s) and saddle point(s) (if any of these exist) of the function

$$z = x^3 + y^3 - 9xy + 27.$$

You may use the second derivative test stated at the beginning of the exam.

1. (6) What is the interval of convergence of the power series $\sum_{n=0}^{\infty} \frac{(x+3)^n}{2^n n!}$?

- A. $(-5, -1]$
- B. $[-5, 1)$
- C. $(-1, 5]$
- D. $[-1, 5)$
- E. $(-\infty, \infty)$

2. (6) What is the value of the improper integral $\int_e^{\infty} \frac{dx}{x(\ln x)^2}$?

- A. -1
- B. 0
- C. $1/e$
- D. 1
- E. the integral diverges

3. (6) What do the level curves of the function $f(x, y) = 3x^2 + 4y^2$ look like?

- A. circles
- B. ellipses
- C. diamonds
- D. parabolas
- E. waves

4. (6) What is the distance from the point $(2, 3, 4)$ to the x -axis?

- A. 2
- B. 3
- C. $\sqrt{13}$
- D. 4
- E. 5

5. (6) Suppose $|\mathbf{a}| = 2$, $|\mathbf{b}| = 3$, and $\mathbf{a} \times \mathbf{b} = \mathbf{0}$. What is $\mathbf{a} \cdot \mathbf{b}$?

A. 0

B. 2

C. 3

D. 6

E. not enough information to decide

6. (6) Suppose that $\frac{n+7}{2-4n} \leq a_n \leq \frac{n^2-n}{(2n+3)^2}$ for all n . What is $\lim_{n \rightarrow \infty} a_n$?

A. $-1/4$

B. 0

C. $1/4$

D. $1/2$

E. not enough information to decide

7. (6) If (a, b) is a critical point of the function $f(x, y)$ and $f_{xx}(a, b) > 0$, $f_{xy}(a, b) = 0$, and $f_{yy}(a, b) > 0$, what kind of critical point is (a, b) ? (You may refer to the Second Derivative Test at the beginning of the other test booklet.)

- A. local minimum
- B. local maximum
- C. saddle point
- D. not enough information to decide

8. (6) If (a, b) is a critical point of the function $f(x, y)$ and $f_{xx}(a, b) > 0$, $f_{xy}(a, b) > 0$, and $f_{yy}(a, b) > 0$, what kind of critical point is (a, b) ? (You may refer to the Second Derivative Test at the beginning of the other test booklet.)

- A. local minimum
- B. local maximum
- C. saddle point
- D. not enough information to decide

9. (6) Consider the series $\sum_{n=10}^{\infty} \frac{1}{n^{\ln \ln n}}$. Which of the following arguments is correct?

- A. this series diverges by the Test for Divergence
- B. this series diverges by comparison to $\sum \frac{1}{n}$
- C. this series converges by comparison to $\sum \frac{1}{n}$
- D. this series diverges by comparison to $\sum \frac{1}{n^2}$
- E. this series converges by comparison to $\sum \frac{1}{n^2}$

10. (6) Suppose that $\nabla f = \langle 2xy + 3x^2y^2, x^2 + 2x^3y \rangle$ and $f(1, 2) = 3$. What is the tangent plane to the surface $z = f(x, y)$ at the point $(1, 2)$?

- A. $z - 3 = (x - 1)/16 + (y - 2)/5$
- B. $z - 3 = 16(x - 1) + 5(y - 2)$
- C. $z - 3 = \langle 16, 5 \rangle t + \langle 1, 2 \rangle$
- D. $z - 3 = (x - 16) + (y - 5)/2$
- E. $z - 3 = (x - 16) + 2(y - 5)$

11. (6) Suppose $\text{proj}_{\mathbf{b}}\mathbf{a} = 3\mathbf{i} - 4\mathbf{j}$ and that the angle between \mathbf{a} and \mathbf{b} is obtuse. What is $\text{comp}_{\mathbf{b}}\mathbf{a}$?

A. 4

B. 5

C. -5

D. $4\mathbf{i} - 3\mathbf{j}$

E. $-3\mathbf{i} + 4\mathbf{j}$

12. (6) Which of the following vectors is parallel to the plane $2x - 3y + 4z = 10$?

A. $\langle 1, -1, 1 \rangle$

B. $\langle 4, -6, 8 \rangle$

C. $\langle 6, 2, 1 \rangle$

D. $\langle 3, 2, 0 \rangle$

E. $\langle -1, -1, 2 \rangle$

13. (6) Suppose that $f(x) = \sum_{n=0}^{\infty} \frac{n^2(x-3)^n}{2^n}$ for $|x-3| < 2$. What is $f^{(38)}(3)$? (The 38th derivative of f at 3.)

A. $\frac{38^2}{2^{38} 38!}$

B. $\frac{38^2}{2^{38}}$

C. $\frac{38^2 38!}{2^{38}}$

D. $\frac{38}{2^{38}}$

E. $\frac{38!}{2^{38}}$

14. (6) What is the domain of the function $\frac{3x^2 + 2y^3}{\sqrt{x^2 + y^2 - 4}}$?

A. the inside of a circle (including the circle)

B. the inside of a circle (not including the circle)

C. the outside of a circle (including the circle)

D. the outside of a circle (not including the circle)

E. the entire plane, except a circle

15. (6) Suppose that $\nabla f = \langle 2xe^{2y}, 2x^2e^{2y} + 3y^2 \rangle$. In which of these directions is $D_{\mathbf{u}}f$ maximized at the point $(2, 0)$?

A. the x direction

B. the y direction

C. $\langle -4, -8 \rangle$

D. $\langle 1/\sqrt{5}, 2/\sqrt{5} \rangle$

E. $\langle -1/\sqrt{5}, -2/\sqrt{5} \rangle$