

Exponential Growth and Decay

10/31/2005

- Derivatives are functions that measure rates of change.
- A rate of change can be a powerful tool for expressing quantitatively a qualitative description.

- We know that a nation's population grows or declines depending on the birth and death rates.
- Does it make sense to say that at any time t , the rate of change of the size of a growing population is proportional to its size?
- Let $y(t)$ be the size of the population at time t .

$$\frac{dy}{dx} = ky; y(0) = y_0.$$

- Note that if $k > 0$, then the population is growing, and if $k < 0$, then the population is decreasing.

Theorem. *The IVP $\frac{dy}{dt} = ky$, $y(0) = y_0$, k constant, has unique solution $y = y_0 e^{kt}$.*

Example

Suppose a bacteria culture grows at a rate proportional to the number of cells present. If the culture contains 700 cells initially and 900 after 12 hours, how many will be present after 24 hours?

Example

Suppose a bacteria culture grows at a rate proportional to the number of cells present. If the culture contains 700 cells initially and 900 after 12 hours, how many will be present after 24 hours?

$$y(12) = 700e^{12k}$$

$$900 = 700e^{12k}$$

$$\frac{900}{700} = e^{12k}$$

$$\ln\left(\frac{900}{700}\right) = 12k$$

$$k = \frac{\ln 900 - \ln 700}{12}$$

Doubling Time and Half-Life

- In an exponential growth model, the doubling time is the length of time required for the population to double.
- In a decay model, the half-life is the length of time required for the population to be reduced to half its size.
- A characteristic of exponential models is that these numbers are independent of the point in time from which the measurement begins.

Example

- A radioactive substance that decays according to an exponential model has a half-life of 600 years. What percentage of an original sample is left after 10 years?

Example

- A radioactive substance that decays according to an exponential model has a half-life of 600 years. What percentage of an original sample is left after 10 years?
- The amount present at time t is given by

$$y(t) = y_0 e^{600k}.$$

Example

- A radioactive substance that decays according to an exponential model has a half-life of 600 years. What percentage of an original sample is left after 10 years?
- The amount present at time t is given by

$$y(t) = y_0 e^{600k}.$$

- To find k we use:

$$\frac{y_0}{2} = y_0 e^{600k}$$

Newton's Law of Cooling

- This law states that a hot object introduced into an environment maintained at a fixed cooler temperature will cool at a rate proportional to the difference between its own temperature and that of the surrounding environment.
- That is, if $y(t)$ is the temperature of the object t units of time after it is introduced into a medium at fixed temperature T_m , we have

$$\frac{dy}{dt} = k(y - T_m); y(0) = y_0$$

where k is a constant.

Example

- Suppose a metal object at 112 degrees Fahrenheit is removed from boiling water and placed on a plate in a room maintained at 68 degrees F.
- Suppose the object cools to 90 degrees in 5 minutes.
- How long will it take to cool to 80 degrees? Note that in this problem, $T_m = 68$, and $y_0 = 112$.

Example

- Suppose a metal object at 112 degrees Fahrenheit is removed from boiling water and placed on a plate in a room maintained at 68 degrees F.
- Suppose the object cools to 90 degrees in 5 minutes.
- How long will it take to cool to 80 degrees? Note that in this problem, $T_m = 68$, and $y_0 = 112$.
- The solution is

$$y - T_m = (y_0 - T_m)e^{kt}$$

$$k = (\ln 22 - \ln 44)/5.$$

Separable Differential Equations

- A first-order differential equation in x and y is called *separable* if it is of the form

$$\frac{dy}{dx} = g(x)h(y).$$

- That is the x 's and dx 's can be put on one side of the equation and the y 's and dy 's on the other

$$\frac{1}{h(y)} dy = g(x) dx$$

$$\int \frac{1}{h(y)} dy = \int g(x) dx$$

Example

- As a review, let's again solve the equation

$$\frac{dy}{dx} = ky$$

by the method of separation of variables.

Justification for the Method of Separation of Variables

- We need to show that given the equation

$$\frac{dy}{dx} = g(x)h(y)$$

the antiderivative of $\frac{1}{h(y)}$ as a function of y equals the antiderivative of $g(x)$ as a function of x .

$$\begin{aligned} f'(x) &= g(x)h(f(x)) \\ \frac{f'(x)}{h(f(x))} &= g(x) \end{aligned}$$

- Let $H(y)$ be any antiderivative of $1/h(y)$

$$\begin{aligned}\frac{d}{dx}H(f(x)) &= H'(f(x))f'(x) \\ &= f'(x)\frac{1}{h(f(x))} \\ &= g(x)\end{aligned}$$

- Let $H(y)$ be any antiderivative of $1/h(y)$

$$\begin{aligned}\frac{d}{dx}H(f(x)) &= H'(f(x))f'(x) \\ &= f'(x)\frac{1}{h(f(x))} \\ &= g(x)\end{aligned}$$

- Thus, the solution $y = f(x)$ satisfies the equation

$$H(f(x)) = \int g(x)dx$$

Examples

- Solve the differential equation

$$\frac{dy}{dx} = \frac{x}{y}.$$


Examples

- Solve the differential equation

$$\frac{dy}{dx} = \frac{x}{y}.$$

- The solution is

$$y^2 - x^2 = 1$$



- Solve the IVP


$$\frac{dy}{dx} = x^2 y^3; y(3) = 1.$$

- Solve the IVP

$$\frac{dy}{dx} = x^2 y^3; y(3) = 1.$$

- The solution is


$$-\frac{1}{2y^2} = \frac{x^3}{3} + C.$$



- From $y(3) = 1$, we find the particular solution:

$$C = -\frac{19}{2}$$

$$y = \sqrt{\frac{1}{19 - \frac{x^3}{3}}}$$



Example

- Solve

$$\frac{dy}{dx} = \frac{2y}{x}.$$


Example

- Solve

$$\frac{dy}{dx} = \frac{2y}{x}.$$

- The solution is

$$y = C_1 x^2$$



Example

- Solve

$$\frac{dy}{dx} = -\frac{x}{2y}.$$


Example

- Solve

$$\frac{dy}{dx} = -\frac{x}{2y}.$$

- The solution is

$$2y^2 + x^2 = C_1.$$



Torricelli's equation

- We found that the equation is of the form

$$y' = k\sqrt{y},$$

where k is a constant.

Torricelli's equation

- We found that the equation is of the form

$$y' = k\sqrt{y},$$

where k is a constant.

- The general form of the solution is

$$y = \left(\frac{1}{2}kx + C_1\right)^2.$$