

Extremal problems consider the minimum and maximum numbers some statistics on a class of graphs can reach. We introduce some of the types of proofs useful in graph theory: Algorithmic, and by construction.

First example

In any simple graph (V, E) , the maximum number of edges is $\binom{|V|}{2}$ *

Proof

In a simple graph, there can be at most one edge per pair of distinct vertices. The maximum number of edges appear in $K_{|V|}$.

This is an extremal problem, since we are looking at the maximum number of edges. The class of graphs here is all simple graphs.

Example

In a bipartite graph with independent sets of size k and m , there can be at most km edges.



Independent sets of size 2 and 4,
8 edges at maximum. km is the number
of edges of $K_{m,k}$

Edges in connected graph

Proposition

The minimum number of edges in a connected graph with n vertices is $n-1$.

Proof

We need to prove two things:

- If a graph with n vertices has fewer than $n-1$ edges, it is not connected.
- There exists a connected graph with n vertices and $n-1$ edges.

(2)

Recall from last week (Monday), that a graph with n vertices and m edges has at least $n-m$ components. Hence, if $m < n-1$, the graph has at least 2 components and is not connected.

Also, the path with n vertices has $n-1$ edges and is connected, proving the minimum is realized.



■

Remark (on the proof technique)

When giving the solution to an extremal problem, there are two parts to be proven:

- That the value we give is minimal (or maximal), i.e. that you cannot give a lower (respectively, higher) value.
- That this value can be realized on at least one graph of the class we consider.

Proposition

Let G be a simple graph with n vertices. If the minimum degree is $\delta(G) \geq \frac{n-1}{2}$, G is connected.

Proof

The minimum degree of the graph means that every vertex should have at least this number of neighbors, in a simple graph.

To prove that G is connected, we must show that there is a path between any pair of vertices $\{u, v\}$. We will in fact prove that there exists a path of length at most 2.

- If $\{u, v\}$ are adjacent, they are obviously in the same component.
- Otherwise, they share at least one neighbor w : There are $n-2$ other vertices, and the sum of their degree is $d(u) + d(v) > n-1$. Hence, $u-w-v$ is a path connecting them.

■

A bound is said to be sharp if improving it (reducing a lower bound or increasing an upper bound) would make the statement wrong.

The bound in the last problem is sharp. To prove it, we give an example of a graph with n vertices and minimum degree $\lfloor \frac{n}{2} \rfloor - 1$ that is not connected: This graph is the disjoint union of $K_{\lfloor \frac{n}{2} \rfloor}$ and $K_{\lceil \frac{n}{2} \rceil}$.



K_5 , degree 4



K_6 , degree 5

11 vertices

Minimum degree is 4, just under $5 = (11-1)/2$.

Graph is disconnected.

Bipartite subgraph


Here we prove that, given a graph G , we can always find a bipartite subgraph with at least a fixed number of edges. We give an algorithmic proof to construct the graph, but a proof can also be done by induction.

Theorem


Every loopless graph $G=(V,E)$ has a bipartite subgraph with at least $|E|/2$ edges.

Proof (algorithmic)

We start with any partition of the vertices into two sets X and Y . Let H be the subgraph containing all the vertices, but only the edges with one endpoint in X and one in Y .



If H has fewer than half the edges incident to a vertex v of X , then it means that v has (in G) more neighbors in X than in Y . To increase the number of edges in H , switch v to Y . The number of edges just increased.




As long as H does not have at least half the edges of G at every vertex, repeat this process. When it terminates, the number of edges in H is always at least half the number of edges of G .

Triangle-free graphs

A graph is said to be triangle-free if it has no three vertices that are all adjacent. In general, a graph G is H -free if it does not contain H as a subgraph.

The Petersen graph is triangle-free (but not bipartite).



Theorem (Mantel, 1907)

The maximum number of edges in a simple triangle-free graph with n vertices is $\lfloor \frac{n^2}{4} \rfloor$

Proof

For the proof, we again need to prove two things:


- that if a graph with n vertices has more than $\lfloor \frac{n^2}{4} \rfloor$ edges, it must have at least one triangle.
- that there always exists a graph with n vertices and $\lfloor \frac{n^2}{4} \rfloor$ edges that has no triangle.

For the first part, take a vertex v of maximal degree Δ . Its Δ neighbors cannot have edges among them.

So every edge of G must have at least one endpoint in a non-neighbor of v , or in v itself.

We give an upper bound on the number of edges:

Since v has maximum degree, the number of edges is at most $\Delta(n-\Delta)$ (because $n-\Delta$ is the number of vertices not adjacent to v). Maximizing $\Delta(n-\Delta)$ gives $\Delta=n/2$. Hence, the number of edges is at most $\lfloor \frac{n^2}{4} \rfloor$



For the second part, we must prove that a triangle-free graph has $\lfloor \frac{n^2}{4} \rfloor$ edges. This is the case of $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$.



We can split 7 vertices into two sets of 3 and 4 vertices, which leads to 12 edges: the smallest integer below $49/4$.

Reference: Douglas B. West. Introduction to graph theory, 2nd edition, 2001. Section 1.3