Math 38 - Graph Theory
Coloring of planar graphs and maps

Nadia Lafrenière 06/02/2020

One of the most interesting theorems in both graph theory and math history is the Four Color Theorem. We see two alternative versions of it, and the history of that theorem.

Six Color Theorem

Theorem

Every planar graph can be colored using six colors.

Lemma

If every subgraph of a graph G has a vertex of degree at most k, then G is (k+1)-colorable.

Proof of the lemma

By induction on the number n of vertices.

- The graph with one vertex is one colorable, and the only vertex has degree o.
- Assume that every graph with at most n vertices such that all its subgraphs contain a vertex of degree at most k is (k+1)-colorable.
- We now look at a graph G with n+1 vertices, and we assume that every subgraph has a vertex of degree at most k. Let v be a vertex in G with degree at most k. Then, the induced subgraph G-v has n vertices and satisfies the induction hypothesis; it is (k+1)-colorable. Since v has k neighbors, it can be colored with one of the k+1 colors.

Proof of the Six Color Theorem

Recall that a planar graph with n vertices has at most 3n-6 edges. That means that there is a vertex with degree at most 5 in every planar graph (and its subgraphs), because otherwise there would be at least 3n edges.

So every subgraph of G is planar and therefore contains a vertex of \bigcirc degree at most 5; using the lemma, it is 6-colorable.

Five Color Theorem

Five Color Theorem (Heawood, 1890) Every planar graph is 5-colorable.

Sketch of proof (details in the textbook)

The proof uses strong induction on the number of vertices.

Base cases: If there are at most 5 vertices, the graph is 5-colorable. Induction step: The graph has at least 6 vertices (and is planar). Using the same argument as for the Six Color Theorem, there is a vertex v of degree at most 5. Also, 6-v satisfies the induction hypothesis, so it is 5-colorable. If v has fewer than 5 neighbors, or if two of the five neighbors of 6 have the same color, we can color v with the fifth color, and we are good.

So assume, by contradiction, that 6-v is 5-colorable, but 6 is not. Necessarily, that means v has neighbors with all different colors: label the five neighbors a,b,c,d and e, with colors respectively 1,2,3,4 and 5. Also, since 6 is not 5-colorable, there is no way we can switch the colors of the neighbors of v to repeat the colors.

That means that, for any pair of colors i and j, the vertices in the neighborhood of 6 that have colors i and j are joined by a path alternating between vertices of color i and color j. Necessarily, these paths must cross at one point, so the graph is not planar (details of the last argument are in the textbook).

The paths must cross at a vertex, because the graph is planar, but this vertex should have a color among 1 and 3 and among 2 and 4; a contradiction. So if G-v is 5-colorable, G must also be 5-colorable.

Path with vertices of color 1 and 3

What color for this vertex?


Path with vertices of color 2 and 4

Remark

Maybe you are (sort of) convinced by this argument that if G-v is five-colorable, then so is G, but what about if V had only 4 neighbors and G-v was 4-colorable? Could we say that G is 4-colorable as well?


Four Color Theorem

A map is a plane drawing. The dual graph of a map is a planar graph.



A coloring of a map is a coloring of its dual graph: two adjacent regions cannot be colored with the same color.

Theorem (Appel, Haken and their computer, 1976)
Every planar graph is 4-colorable. Every map is 4-colorable.



A bit of history on the Four Color Theorem

1852: Earliest known posing of the conjecture (in terms of map coloring)

1878: Popularization of the conjecture at a mathematical meeting

1879: First "proof", similar to the proof of the Five Color Theorem

1879-1976: Plenty of incomplete proofs

1976: Kenneth Appel and Wolfgang Haken prove the theorem using computer enumeration. They reduced the general case to 1,834 cases that the computer has to check. It takes over 1,000 hours to the computer to complete the proof, which cannot be read by a human being.

1996: Simplification of the proof to (only) 633 cases

2005: Computer verification of the proof

2020+? Can we imagine a proof that would be readable by a human?

Reference: Douglas B. West. Introduction to graph theory, 2nd edition, 2001. Section 6.3