

Number Theory Web Seminar
March 12, 2026

The Erdős–Straus Conjecture


Carl Pomerance, Dartmouth College

Joint work with


Andreas Weingartner, Southern Utah University



Andreas Weingartner



Paul Erdős



Ernst Straus

(Drawing by LeUyen Pham, illustrator of *The Boy Who Loved Math*, by Deborah Heiligman)

In 1948, Paul Erdős and Ernst Straus conjectured that for every integer $n \geq 2$, there are positive integers x, y, z such that

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

The first question: **Why make this conjecture???**



The Rhind papyrus, ca. 1500 BCE

Apparently the ancient Egyptians were especially fond of fractions with numerator 1, so-called *unit fractions*. To represent other fractions, they would find some unit fractions that summed to what they wanted.

For example, consider $5/7$. We have

$$\frac{5}{7} = \frac{1}{2} + \frac{1}{5} + \frac{1}{70}.$$

The Rhind papyrus gave a list of such representations, now called Egyptian fractions.

Note that the above decomposition of $5/7$ was found by the “greedy algorithm”, where each fraction is chosen as large as possible without exceeding the target. For example, $1/2 < 5/7 < 1/1$, so we choose $1/2$. We have $5/7 - 1/2 = 3/14$ and $1/5 < 3/14 < 1/4$, so we choose $1/5$. What’s left is $1/70$, and we’re done.

The greedy algorithm for Egyptian fractions was first described by Fibonacci about 800 years ago.

Note that the greedy algorithm does not always give the shortest representation. Lets try $4/17$. The greedy algorithm gets

$$\frac{4}{17} = \frac{1}{5} + \frac{1}{29} + \frac{1}{1233} + \frac{1}{3039345},$$

but

$$\frac{4}{17} = \frac{1}{6} + \frac{1}{17} + \frac{1}{102}$$

is simpler.

One might try and describe the set of rationals which have a shortest representation as a sum of k unit fractions.

When $k = 1$, we have the unit fractions themselves.

When $k = 2$, we have the identity

$$\frac{2}{n} = \frac{1}{n} + \frac{1}{n},$$

which shows that each $2/n$ is in class 2 for n odd. But there are many more fractions in the class 2, for example $5/6$ is.

Theorem (Stewart, 1964). If $(m, n) = 1$, we have m/n the sum of 2 unit fractions if and only if m is a divisor of the sum of two coprime divisors of n .

For example, 2 and 3 are coprime divisors of 6 and 5 is a divisor of $2 + 3$, so $5/6$ is the sum of 2 unit fractions. But $5/7$ is not, nor is $4/17$, nor is any m/p with p an odd prime and $m \nmid p + 1$.

Proof. Suppose $a, b \mid n$ with $(a, b) = 1$ and $m \mid a + b$. Write $n = abc$ and $a + b = md$. Then

$$\frac{m}{n} = \frac{md}{nd} = \frac{a + b}{abcd} = \frac{1}{bcd} + \frac{1}{acd}.$$

Theorem (Stewart, 1964). If $(m, n) = 1$, we have m/n the sum of 2 unit fractions if and only if m is a divisor of the sum of two coprime divisors of n .

Proof, cont'd. Now suppose that $m/n = 1/u + 1/v$. Say $(u, v) = c$, $u = ac$, $v = bc$. Then

$$\frac{m}{n} = \frac{1}{u} + \frac{1}{v} = \frac{a+b}{abc}.$$

Let $d = (a+b, c)$. Since $(a, b) = 1$ we thus have $md = a+b$, $nd = abc$. So a, b are coprime divisors of n , and we're done.

A corollary of Stewart's theorem is that for each fixed m , almost all n have m/n the sum of two unit fractions! Indeed, if n is divisible by some prime $p \equiv -1 \pmod{m}$, then since $p, 1$ are coprime divisors of n and $m \mid p+1$, we have m/n the sum of two unit fractions. And the number of integers $n \leq x$ **not** divisible by any prime $p \equiv -1 \pmod{m}$ is $O_m(x/(\log x)^{1/\varphi(m)})$. Here $\varphi(m)$ is Euler's function.

This count can usually be improved. For example, if $m = 7$, then $7/n$ is not the sum of two unit fractions when n avoids primes that are $\equiv 3, 5, \text{ or } 6 \pmod{7}$ and we get a count of the shape $x/(\log x)^{1/2+o(1)}$.

More generally, If m/n is not the sum of two unit fractions, then essentially n takes only primes in a subgroup of $(\mathbf{Z}/m\mathbf{Z})^*$ that avoids -1 . (This result is due to Huang and Vaughan, 2011.)

So, the situation when m/n is the sum of two unit fractions is basically understood.

Which brings us to the sum of three unit fractions:

$$\frac{m}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

Note that if $m \leq 3$, then the problem is trivial.

So, the case of $m = 4$, the arena of the Erdős–Straus conjecture, is the first interesting case.

But why would one suspect that for every $n \geq 2$ with $m = 4$ there is a solution? Could this really be true?

First note that if each $4/p$ with p prime is a sum of three unit fractions, then so is each $4/n$. Indeed, if $p \mid n$, say $n = jp$, then dividing a representation for $4/p$ by j gets a representation for $4/n$.

Further, if $p \equiv 3 \pmod{4}$, then $4 \mid p+1$ so $4/p$ is the sum of two unit fractions (therefore also the sum of three).

Suppose $p+1$ is divisible by a prime $q \equiv 3 \pmod{4}$, so write $q = 4k - 1$ and $p+1 = jq$. Then $p = j(4k - 1) - 1 = 4jk - j - 1$ and

$$\frac{4}{p} = \frac{4jk}{pjk} = \frac{1}{jk} + \frac{j+1}{pjk} = \frac{1}{jk} + \frac{1}{pk} + \frac{1}{pjk}.$$

This is an early result of Obláth and it immediately implies that the number of exceptional $p \leq x$ for which the Erdős–Straus conjecture might be false is $O(x/(\log x)^{3/2})$.

This generalizes as follows: Again we have primes p, q with $q \equiv 3 \pmod{4}$. Write $q = 4uvw - 1$ and assume that p satisfies $pv \equiv -u \pmod{q}$. (The Obláth case had $u = v = 1$.) Then there is some integer t with $pv + u = tq = 4tuvw - t$, that is,

$$4tuvw = pv + u + t.$$

Dividing this equation by $ptuvw$, we get

$$\frac{4}{p} = \frac{1}{tuw} + \frac{1}{ptvw} + \frac{1}{puvw}.$$

This kind of thing can go in two directions:


- (1) get congruences for p where we know the conjecture holds,
- (2) get so many congruences that there are only a few p 's left in doubt.

The first is an excellent aid for an exhaustive search for counterexamples.

Using congruences, as reported in Mordell's famous book on Diophantine equations, one learns that the Erdős–Straus conjecture holds for every prime p except possibly for the quadratic residues mod 840, that is, except for those p with

$$p \equiv 1, 121, 169, 289, 361, \text{ or } 529 \pmod{840}.$$

Using congruences such as these, about 12 years ago, Salez verified the Erdős–Straus conjecture to 10^{17} . Just recently, Mihnea and Dumitru extended the search to 10^{18} .



Robert C. Vaughan


In 1970 Vaughan used the second approach and the large sieve to get an excellent upper bound for the number of possible exceptions up to N : It is $O(N/\exp(c(\log N)^{2/3}))$ for an appropriate positive constant c .

If you can't prove it, generalize it. . .



Wacław Sierpiński

Sierpiński conjectured that every $5/n$ for $n \geq 2$ is a sum of three unit fractions. (With Weingartner, we have verified this to 10^{18} .)



Andrzej Schinzel

Schinzel then generalized further: For every $m \geq 4$, we have m/n the sum of three unit fractions for all $n > N_m$, a constant depending on m . With Weingartner, we have investigated this numerically for m up to 15. For example, when $m = 8$ we have checked up to 10^{13} and the only exceptional n found are 1, 2, 3, 11, 17, 131, 241. See the table on the next slide.

m	all exceptions $n \leq N$	Count	N
4	1	1	10^{18}
5	1	1	10^{18}
6	1	1	10^{13}
7	1, 2	2	10^{13}
8	1, 2, 3, 11, 17, 131, 241	7	10^{13}
9	1, 2, 5, 11, 19	5	10^{12}
10	1, 2, 3, 7, 11, 43, 61, 67, 181	9	10^{12}
11	1, 2, 3, 4, 37	5	10^{12}
12	1, 2, 3, 5, 7, 13, 25, 29, 31, 37, 73, 97, 193, 433, 577, 1129, 1657, 1873, 2521, 2593, 3433, 10369, 12049, 12241	24	10^{12}
13	1, 2, 3, 4, 5, 7, 14, 53, 61, 67, 79, 211, 281	13	10^{12}
14	1, 2, 3, 4, 5, 17, 19, 29, 59, 257, 353, 841	12	10^{12}
15	1, 2, 3, 4, 8, 16, 17, 19, 23, 31, 34, 47, 53, 61, 79, 113, 122, 137, 151, 197, 226, 233, 271, 541, 1103, 1171, 1367, 4201, 6301, 12601, 16831, 20521	32	10^{12}

Already in his 1970 paper, Vaughan proved a general upper bound for exceptions to the Schinzel conjecture: The number of $n \leq N$ for which m/n is not the sum of three unit fractions is $O(N/\exp(c(m)(\log N)^{2/3}))$, where $c(m) > 0$.

With Weingartner, we were able to prove this with $c(m) = c/\varphi(m)^{1/3}$, with c an absolute positive constant, uniformly for $m \leq (\log N)^2$.

But mainly my work with Weingartner deals with the exceptional set in the Schinzel variant. Schinzel's conjecture is that for each m there is some N_m such that when $n > N_m$ we have m/n the sum of three unit fractions. How large is this N_m ? For example, the Erdős–Straus conjecture is that $N_4 = 1$. And empirically it seems that $N_8 = 241$. We show that as m gets large, exceptions become enormous.

Theorem (Pomerance & Weingartner). For each $\epsilon > 0$ there is a bound m_ϵ such that if $m > m_\epsilon$ there is a number $n > \exp(m^{1/3-\epsilon})$ with m/n not the sum of three unit fractions.

In fact, with $N = \exp(m^{1/3-\epsilon})$, we show that most primes p in $(N, 2N]$ have m/p not the sum of three unit fractions.

Our proof uses many ideas from a recent paper of Elsholtz and Tao on counting the number of triples x, y, z where $4/n = 1/x + 1/y + 1/z$. They also do a good job of citing the many researchers who have obtained partial results.



Christian Elsholtz



Paul Erdős & Terence Tao

Let $m \geq 4$ and let p be a prime. The first observation is that solutions to $m/p = 1/x + 1/y + 1/z$ come in two types. A Type I solution has p dividing just one of x, y, z , while a Type II solution has p dividing two of x, y, z . (Since the smallest of x, y, z must be $< p$, it follows that p cannot divide all of x, y, z .)

The next observation is that it is possible to give parametrizations of the two types.

There is a Type I solution if and only if there are positive integers a, d, f such that

$$f \mid ma^2d + 1, \quad mad \mid p + f.$$

There is a Type II solution if and only if there are positive integers a, b, e with

$$e \mid a + b, \quad mab \mid p + e.$$

Lets focus on Type I solutions:

$$f \mid ma^2d + 1, \quad mad \mid p + f.$$

Let N be a large function of m , say about $\exp(m^{1/3})$, to get a feel for the argument. We try to count primes $p \in (N/2, N]$ for which a Type I solution exists. We see that p is in a residue class modulo mad , so a ready tool to use is the Brun–Titchmarsh theorem. This gives the bound

$$\ll \sum_{a,d: ad < 3N/m} \frac{N\tau(ma^2d + 1)}{\varphi(mad) \log(2 + N/mad)}.$$

Then using a calculation in Elsholtz–Tao for part of the range, and another argument for another part, we find that the number of primes $p \in (N/2, N]$ with a Type I solution is

$$\ll \frac{N}{\varphi(m)} (\log N)^2 (\log m)^2.$$

The trick is to then choose the relationship between N, m so that this is $< \epsilon N / \log N$, which will hold if

$$N \leq \exp((\varphi(m)/C \log^2 m)^{1/3})$$

for C large. With this choice, we have most primes in $(N/2, N]$ do not have a Type I solution.

So, it then comes down to showing that most primes in this range also do not have a Type II solution. Similar methods show that there are fewer primes of this type, so we conclude that most primes p in $(N/2, N]$ with N bounded as above have m/p not the sum of three unit fractions.

One of the theorems mentioned in this talk says that in a certain range there are few exceptions, and in another range there are few solutions. Lets compare these ranges.

We've just seen that near $\exp(m^{1/3})$ or smaller most primes p will not have m/p the sum of three unit fractions.

Earlier we saw that for $m \leq \log^2 N$ most primes p near N will have solutions. This translates to N near $\exp(m^{1/2})$ or larger.

This raises the question of where the transition is from almost never to almost always to always. Our proofs suggest that the average count of solutions for a given p is about $(\log p)^3/m$. We follow the thoughts of Elsholtz–Tao in the $m = 4$ case that there is a Poisson process at work, with the likelihood of no solution for m/p being about $\exp(-(\log p)^3/m)$. This suggests if $p < \exp(m^{1/3-\epsilon})$ it's unusual to have m/p the sum of three unit fractions. Once p grows to about $\exp(m^{1/3+\epsilon})$ it is now common for there to be a solution but many times there are not. At $\exp(m^{1/2-\epsilon})$ it's the same situation but the exceptions are very sparse. And once $p > \exp(m^{1/2+\epsilon})$, it is always true.

Recall that Schinzel conjectured there is a number N_m such that if $n > N_m$ then m/n is the sum of three unit fractions. If so, there is a perhaps smaller bound N'_m such that m/p is a sum of three unit fractions for all primes $p > N'_m$. The above suggests that N'_m can be taken as $\exp(m^{1/2+o(1)})$.

This says little about the $m = 4$ case where this story began. Feel free to have at it!

For further reading on this subject, see

T. Bloom and C. Elsholtz, Egyptian fractions, *Nieuw Arch. Wisk.* **23** (2022), 237–245.

C. Elsholtz and T. Tao, Counting the number of solutions to the Erdős–Straus equation on unit fractions, *J. Aust. Math. Soc.* **94** (2013), 50–105.

C. Pomerance and A. Weingartner, Exceptions to the Erdős–Straus–Schinzel conjecture, *Ramanujan J.*, to appear.

(These papers are all available on arXiv and the slides for this talk are on my home page.)

Thank You!