

ERRATA:
ON THE ARITHMETIC DIMENSION OF TRIANGLE GROUPS

STEVE NUGENT AND JOHN VOIGHT

This note gives errata for the article *On the arithmetic dimension of triangle groups* [1]. The authors thank Giovanni Panti and Cherng-tiao Perng.

- (1) (1.2): there is a \mathbb{Q} missing, so it should read

$$E = \mathbb{Q}(\mathrm{Tr} \Delta^{(2)}) = \mathbb{Q}(\{\mathrm{Tr}(\delta^2) : \delta \in \Delta\}) = \mathbb{Q}(\lambda_a, \lambda_b, \lambda_c, \lambda_{2a}\lambda_{2b}\lambda_{2c})$$

- (2) (1.3): in two lines before, this is the preimage of $\Delta^{(2)}$, and the isomorphism requires $b \neq \infty$. (When $b = \infty$, necessarily $A \simeq M_2(E)$ because δ_b is parabolic, so A has a zerodivisor.)
- (3) Lemma 1.9: the cardinality signs are missing, so it should read

$$\begin{aligned} \mathrm{adim}(a, b, c) &= \#\{k \in (\mathbb{Z}/2m\mathbb{Z})^\times / H : \sigma_k(\beta) > 0\} \\ &= \#\{k \in (\mathbb{Z}/2m\mathbb{Z})^\times / H : \kappa(a, b, c; k) < 0\}. \end{aligned}$$

- (4) Lemma 3.4: there is a typo in the inequality, and the statement should read: $\kappa(a, b, c; k) \geq 0$ if and only if (3.5) holds. In the proof, we let $f(z) = \kappa(a, b, c; k)$ and considered when $f(z) \geq 0$. The remaining statements using Lemma 3.4 (Corollary 3.6, Proposition 4.6, Theorem 5.2) use the corrected Lemma 3.4; before (3.8) the inequality should be switched to > 0 ; the rest of the paper remains unchanged.
- (5) Proof of Lemma 3.4, “The discriminant simplifies as”: This should read

$$\sqrt{t^2 - 4n} = \sqrt{4 \cos^2 \frac{k_a \pi}{a} \cos^2 \frac{k_b \pi}{b} - 4 \cos^2 \frac{k_a \pi}{a} - 4 \cos^2 \frac{k_b \pi}{b} + 4}$$

(so the final -4 should be $+4$), as shown on the next line.

- (6) Proof of Proposition 4.6, “In particular $1 \leq 3q/a$ ”: Should be “In particular, $1 < 3q/a$.”
- (7) Beginning of section 5: should be “section”, not “chapter”.
- (8) Line 2 of Algorithm 1: Typo, should be $\max(48, 2r)$ (according to Lemma 4.10).
- (9) Line 5 of Algorithm 3: “for” should be **for**, and it is missing **do**. This step just initializes **divisors** to be an array of 1s.
- (10) Line 5 of Algorithm 4: should be $(\mathbf{a}, \mathbf{b}, \mathbf{c}, r)$.
- (11) Theorem 5.2: To be more precise, this theorem proves the correctness of `FIND_ARITHMETIC`.
- (12) Proof of Theorem 5.2: The middle paragraph is a bit muddled. Here is a revised version that hopefully makes the logic clear.

We have from Lemma 3.4 that for every (a, b, c) and every $k \in \mathbb{Z}_{>0}$ with $k \in (\mathbb{Z}/2m\mathbb{Z})^\times$, if $\kappa(a, b, c; k) > 0$ then

$$c < \frac{k_c}{|k_a/a + k_b/b - 1|} \leq \frac{k}{|k_a/a + k_b/b - 1|} = \frac{kab}{|k_a b + k_b a - ab|}.$$

Suppose (a, b, c) is r -arithmetic, and consider the set of primes $q < c/2$ with $q \nmid m$. As in the proof of Lemma 4.3, these primes are distinct in $(\mathbb{Z}/2m\mathbb{Z})^\times/H$, so there at most $r-1$ of them for which $\kappa(a, b, c; q) < 0$; so among any r of them, there is at least one with $\kappa(a, b, c; q) > 0$. Putting these observations together, we find that in any set of r primes with $q \nmid ab$, there is a prime q in the set where at least one of the following holds:

$$\text{either } q \mid c \text{ or } c < 2q \text{ or } c < \left\lceil \frac{qab}{|q_a b + q_b a - ab|} \right\rceil.$$

To finish, suppose (a, b, c) is r -arithmetic, $c > 2*\text{maxNDP}$, and $c > \text{bound}$, where **bound** is in the keyset of **boundToPrimes**. Let

$$B = \{q : q \in \text{boundToPrimes}[\text{bound}'] \text{ and } \text{bound}' \leq \text{bound}\}.$$

Then there exist at most $r-1$ primes $q \in B$ that do not divide c . The algorithm partitions B into r sets, and lets each **divisor** in **divisors** be the product of primes in one such set. Hence, c must be a multiple of at least one **divisor** in **divisors**. Therefore, the algorithm checks all possible r -arithmetic triples (a, b, c) .

REFERENCES

- [1] Steve Nugent and John Voight, *On the arithmetic dimension of triangle groups*, Math. Comp. **86** (2017), no. 306, 1979–2004.