

How Many Borel Sets are There?

Object. *This series of exercises is designed to lead to the conclusion that if $\mathcal{B}_{\mathbf{R}}$ is the σ -algebra of Borel sets in \mathbf{R} , then*

$$\text{Card}(\mathcal{B}_{\mathbf{R}}) = \mathfrak{c} := \text{Card}(\mathbf{R}).$$

This is the conclusion of problem 4. As a bonus, we also get some insight into the “structure” of $\mathcal{B}_{\mathbf{R}}$ via problem 2. This just scratches the surface. If you still have an itch after all this, you want to talk to a set theorist. This treatment is based on the discussion surrounding [1, Proposition 1.23] and [2, Chap. V §10 #31].

For these problems, you will definitely want to have a close look at [1, §0.4] on well ordered sets. Note that by [1, Proposition 0.18], there is an uncountable *well ordered* set Ω such that for all $x \in \Omega$, $I_x := \{y \in \Omega : y < x\}$ is countable. The elements of Ω are called *countable ordinals*. We let $1 := \inf \Omega$. If $x \in \Omega$, then $x + 1 := \inf\{y \in \Omega : y > x\}$ is called the *immediate successor* of x . If there is a $z \in \Omega$ such that $z + 1 = x$, then z is called the *immediate predecessor* of x . If x has no immediate predecessor, then x is called a *limit ordinal*.¹

1. Show that $\text{Card}(\Omega) \leq \mathfrak{c}$. (This follows from [1, Propositions 0.17 and 0.18]. Alternatively, you can use transfinite induction to construct an injective function $f : \Omega \rightarrow \mathbf{R}$.)

2. If X is a set, let $\mathcal{P}(X)$ be the set of subsets of X — i.e., $\mathcal{P}(X)$ is the *power set* of X . Let $\mathcal{E} \subset \mathcal{P}(X)$. The object of this problem is to give a “concrete” description of the σ -algebra $\mathcal{M}(\mathcal{E})$ generated by \mathcal{E} . (Of course, we are aiming at describing the Borel sets in \mathbf{R} which are generated by the collection \mathcal{E} of open intervals.) For convenience, we assume that $\emptyset \in \mathcal{E}$.

Let

$$\mathcal{E}^c := \{E^c : E \in \mathcal{E}\} \quad \text{and} \quad \mathcal{E}_\sigma = \left\{ \bigcup_{i=1}^{\infty} E_i : E_i \in \mathcal{E} \right\}.$$

(Note, I just mean that \mathcal{E}_σ is the set of sets formed from countable unions of elements of \mathcal{E} . Since $\emptyset \in \mathcal{E}$, $\mathcal{E} \subset \mathcal{E}_\sigma$.)

We let $\mathcal{F}_1 := \mathcal{E} \cup \mathcal{E}^c$. If $x \in \Omega$, and if x has an immediate predecessor y , then we set

$$\mathcal{F}_x := (\mathcal{F}_y)_\sigma \cup ((\mathcal{F}_y)_\sigma)^c.$$

¹The set of countable ordinals has a rich structure. We let $2 := 1 + 1$, and so on. The set $\{n \in \mathbf{N}\} \subset \Omega$ is countable, and so has a supremum ω (see [1, Proposition 0.19]). Then there are ordinals $\omega + 1, \omega + 2, \dots, 2\omega, 2\omega + 1, \dots, \omega^2, \omega^2 + 1, \dots, \omega^\omega$, and so on.

If x is a limit ordinal, then we set

$$\mathcal{F}_x := \bigcup_{y < x} \mathcal{F}_y.$$

We aim to show that

$$\mathcal{M}(\mathcal{E}) = \bigcup_{x \in \Omega} \mathcal{F}_x \tag{†}$$

- (a) Observe that $\mathcal{F}_1 \subset \mathcal{M}(\mathcal{E})$.
- (b) Show that if $F_y \subset \mathcal{M}(\mathcal{E})$ for all $y < x$, then $F_x \subset \mathcal{M}(\mathcal{E})$. Then use transfinite induction to conclude that $\mathcal{F}_x \subset \mathcal{M}(\mathcal{E})$ for all $x \in \Omega$.
- (c) Show that the right-hand side of (†) is closed under countable unions.
- (d) Conclude that $\bigcup_{x \in \Omega} \mathcal{F}_x$ is a σ -algebra, and that (†) holds.

3. Recall that if A and B are sets, then $\prod_{a \in A} B$ is simply the set of functions from A to B . For reasons that are unclear to me, this set is usually written B^A . Notice that $\prod_{i=1}^{\infty} B = \prod_{i \in \mathbf{N}} B$ is just the collection of sequences in B . Notice also that $\text{Card}(B^A)$ depends only on $\text{Card}(A)$ and $\text{Card}(B)$.

- (a) Check that

$$\prod_{i=1}^{\infty} \left(\prod_{j=1}^{\infty} B \right) = \prod_{(i,j) \in \mathbf{N} \times \mathbf{N}} B. \tag{*}$$

Thus the cardinality of either side of (*) is the same as $\prod_{i=1}^{\infty} B$.

- (b) Use these observations together with the fact that $\text{Card}(\prod_{i=1}^{\infty} \{0, 1\}) = \mathfrak{c} := \text{Card}(\mathbf{R})$ (which follows from [1, Proposition 0.12]) to show that

$$\text{Card} \left(\prod_{i=1}^{\infty} \mathbf{R} \right) = \mathfrak{c}.$$

- (c) Show that if $\text{Card}(\mathcal{E}) = \mathfrak{c}$, then $\text{Card}(\mathcal{E}_{\sigma}) = \mathfrak{c}$.

4. Let $\mathcal{B}_{\mathbf{R}}$ be the σ -algebra of Borel sets in \mathbf{R} . In [1, Proposition 0.14(b)], it is shown that if $\text{Card}(A) \leq \mathfrak{c}$ and if $\text{Card}(Y_x) \leq \mathfrak{c}$ for all $x \in A$, then $\bigcup_{x \in A} Y_x$ has cardinality bounded by \mathfrak{c} . By following the given steps, use this observation, as well as problems 2 and 3, to show that

$$\text{Card}(\mathcal{B}_{\mathbf{R}}) = \mathfrak{c}. \tag{‡}$$

- (a) Let \mathcal{E} be the collection of open intervals (including the empty set) in \mathbf{R} . Then $\text{Card}(\mathcal{E}) = \mathfrak{c}$.
- (b) $\mathcal{B}_{\mathbf{R}} = \mathcal{M}(\mathcal{E})$.
- (c) Define \mathcal{F}_x as in problem 2. Use transfinite induction and problem 3 to prove that $\text{Card}(F_x) = \mathfrak{c}$ for all $x \in \Omega$.
- (d) Use problem 2 to conclude that $\mathcal{M}(\mathcal{E}) = \mathcal{B}_{\mathbf{R}}$ has the cardinality claimed in (\dagger) .²

References

- [1] Gerald B. Folland, *Real analysis*, Second, John Wiley & Sons Inc., New York, 1999. Modern techniques and their applications, A Wiley-Interscience Publication.
- [2] Anthony W. Knap, *Basic real analysis*, Cornerstones, Birkhäuser Boston Inc., Boston, MA, 2005. Along with a companion volume *Advanced real analysis*.

²It is my *understanding* that the classes \mathcal{F}_x are all distinct; that is, $\mathcal{F}_x \subsetneq \mathcal{F}_y$ if $x < y$ in Ω . But I don't have a reference or a proof at hand.