Winter 2020 Math 126
Topics in Applied Mathematics
Data-driven Uncertainty Quantification

Yoonsang Lee (yoonsang.lee@dartmouth.edu)

Lecture 3: Information Theory

3.1 Entropy

Def. The entropy $H(X)$ of a random variable X with density $p(x)$ is defined as

$$H(X) = -\int_S p(x) \ln p(x) dx,$$

where S is the support of $p(x)$ (that is, the set where $p(x)$ is not zero).

Entropy depends only on the density $p(x)$ and thus entropy is sometime written as $H(p)$ rather than $H(X)$.

3.1 Entropy

Example. Let X is a Gaussian with density $p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{x^2}{2\sigma^2}}$.

$$\begin{aligned} H(p) &= -\int p \ln p dx \\ &= -\int p \left[-\frac{x^2}{2\sigma^2} - \ln \sqrt{2\pi\sigma^2} \right] dx \\ &= \frac{E[X^2]}{2\sigma^2} + \frac{1}{2} \ln 2\pi\sigma^2 \\ &= \frac{1}{2} + \frac{1}{2} \ln 2\pi\sigma^2 \\ &= \frac{1}{2} \ln 2\pi e\sigma^2 \end{aligned} \tag{1}$$

Note. For a n-dimensional Gaussian X with mean zero and covariance K, $H(p) = \frac{1}{2} \ln (2\pi e)^m |K|$ where $|K|$ is the determinant of K.

3.2 Joint and Conditional Entropy

Def. The entropy of a set $X_1, X_2, ..., X_n$ of random variables with density $p(x_1, x_2, ..., x_n)$ is defined as

$$H(p(x_1, x_2, ..., x_n)) = -\int p(x_1, x_2, ..., x_n) \ln p(x_1, x_2, ..., x_n) dx_1 \cdots dx_n.$$

Def. If X and Y have a joint density $p(x, y)$, the conditional entropy $H(X|Y)$ is defined as

$$H(X|Y) = -\int p(x, y) \ln p(x|y) dxdy.$$

3.2 Joint and Conditional Entropy

Def. The entropy of a set $X_1, X_2, ..., X_n$ of random variables with density $p(x_1, x_2, ..., x_n)$ is defined as

$$H(p(x_1, x_2, ..., x_n)) = -\int p(x_1, x_2, ..., x_n) \ln p(x_1, x_2, ..., x_n) dx_1 \cdots dx_n.$$

Def. If X and Y have a joint density $p(x, y)$, the conditional entropy $H(X|Y)$ is defined as

$$H(X|Y) = -\int p(x, y) \ln p(x|y) dxdy.$$

Q. Why not $-\int p(x|y) \ln p(x|y) dxdy$?

3.2 Joint and Conditional Entropy

Def. The entropy of a set $X_1, X_2, ..., X_n$ of random variables with density $p(x_1, x_2, ..., x_n)$ is defined as

$$H(p(x_1, x_2, ..., x_n)) = -\int p(x_1, x_2, ..., x_n) \ln p(x_1, x_2, ..., x_n) dx_1 \cdots dx_n.$$

Def. If X and Y have a joint density $p(x, y)$, the conditional entropy $H(X|Y)$ is defined as

$$H(X|Y) = -\int p(x, y) \ln p(x|y) dxdy.$$

Q. Why not $-\int p(x|y) \ln p(x|y) dxdy$?
Fact. $H(X|Y) = H(X, Y) - H(Y)$

3.3 Relative Entropy and Mutual Information

Def. The relative entropy (or Kullback-Leibler distance) $D(p, g)$ between two densities p and q is defined by

$$D(p, q) = \int p \ln \frac{p}{q} dx$$

D is a measure of the inefficiency of assuming that the distribution is q when the true distribution is p.

Def. The mutual information $I(X, Y)$ between two random variables with joint density $p(x, y)$ is defined as

$$I(X, Y) = \int p(x, y) \ln \frac{p(x, y)}{p(x)p(y)} dxdy.$$

Note.

$I(X, Y) = D(p(x, y), p(x)p(y)) = H(X) + H(Y) - H(X, Y)$.

3.3 Relative Entropy and Mutual Information

Example. Let (X, Y) is a Gaussian with mean $(0, 0)$ and a covariance $K = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$.

3.3 Relative Entropy and Mutual Information

Example. Let (X, Y) is a Gaussian with mean $(0, 0)$ and a covariance $K = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$.
$H(X) = H(Y) = \frac{1}{2}\ln(2\pi e)$ and $H(X, Y) = \frac{1}{2}\ln(2\pi e)^2(1 - \rho^2)$.
Therefore $I(X, Y) = H(X) + H(Y) - H(X, Y) = -\frac{1}{2}\ln(1 - \rho^2)$. If $\rho = 0$, X and Y are independent and the mutual information is 0.
If $\rho = \pm 1$, X and Y are perfectly correlated and the mutual information is infinite.
Note. X and Y are Gaussian and thus zero correlation implies independence.

3.4 Properties of entropy, relative entropy, and mutual information

Theorem.

$$D(p,q) \geq 0$$

with equality iff $p = q$ almost everywhere.

Proof.

$$\begin{aligned} -D(p,q) &= \int p \ln \frac{q}{p} dx \\ &\leq \ln \int p \frac{q}{p} dx \quad \text{from Jensen's inequality} \\ &= \ln \int g \\ &\leq \ln 1 = 0. \end{aligned} \tag{2}$$

Corollary. $I(X,Y) \geq 0$ with equality iff X and Y are independent.

3.4 Properties of entropy, relative entropy, and mutual information

Corollary. $H(X|Y) \leq H(X)$ with equality iff X and Y are independent.

3.4 Properties of entropy, relative entropy, and mutual information

Corollary. $H(X|Y) \leq H(X)$ with equality iff X and Y are independent. That is, collecting data decreases uncertainty (yay!).

3.4 Properties of entropy, relative entropy, and mutual information

Theorem. (Chain rule for entropy)

$$H(X_1, X_2, ..., X_n) = \sum H(X_i | X_1, X_2, ..., X_{i-1}).$$

Proof. Homework.

Corollary.

$$H(X_1, X_2, ..., X_n) \leq \sum H(X_i)$$

Hadamard's inequality. If X is a Gaussian distribution with mean 0 and a covariance K, we have

$$|K| \leq \Pi_{i=1}^{n} K_{ii}$$

where $|K|$ is the determinant of K.

3.4 Properties of entropy, relative entropy, and mutual information

In Lecture 1, we have seen that the probability density maximizing entropy with a given mean and a variance is Gaussian. Now we show the following general result.

Theorem. Let the random vector $X \in \mathbb{R}^n$ have zero mean and covariance K. Then

$$H(X) \leq \frac{1}{2} \ln(2\pi e)^n |K|,$$

with equality iff X is Gaussian is the covariance K and mean zero. $|K|$ is the determinant of K.

Proof. Let $g(x)$ be any density satisfying $\int g(x) x_i x_j dx_i dx_j = K_{ij}$ for all i, j. Let ϕ_K be the density of the Gaussian $N(0, K)$. Then

$$\begin{aligned} 0 &\leq D(g, \phi_K) \\ &= \int g \ln(g/\phi_K) \\ &= -h(g) - \int g \ln \phi_K \\ &= -h(g) - \int \phi_K \ln \phi_K \\ &= -h(g) + h(\phi_K). \end{aligned} \tag{3}$$

3.4 Properties of entropy, relative entropy, and mutual information

Theorem. (Estimation error) For any one-dimensional random variable X and estimator $\hat{X}$,

$$E[(X-\hat{X})^2] \geq \frac{1}{2\pi e} e^{2H(X)},$$

with equality iff X is Gaussian and $\hat{X}$ is the mean of X.

Proof. Let $\hat{X}$ be any estimator of X. Then

$$\begin{aligned} E[(X-\hat{X})^2] &\geq \min_{\hat{X}} E[(X-\hat{X})^2] \\ &= E[(X-E[X])^2] \\ &= Var(X) \\ &\geq \tfrac{1}{2\pi e} e^{2H(X)}. \end{aligned} \tag{4}$$

Homework

- Draw n values of the standard normal random variable, X.
- When $Y = X^2$, calculate $D(X, Y)$ using the sample. If you use a histogram in a sense, change the number of bins and check the change of the relative entropy.
- Compare the relative entropy with an analytic solution.