


Central Limit Theorem

Bernoulli Trials

11/01/2005

Continuous Probability Densities

- Let us construct a spinner, which consists of a circle of unit circumference and a pointer.



- The experiment consists of spinning the pointer and recording the label of the point at the tip of the pointer.

- We let the random variable X denote the value of this outcome.
- The sample space is clearly the interval $[0, 1)$.
- It is necessary to assign the probability 0 to each outcome.
- The probability
$$P(0 \leq X \leq 1)$$
should be equal to 1.

- We would like the equation

$$P(c \leq X < d) = d - c$$

to be true for every choice of c and d .

- If we let $E = [c, d]$, then we can write the above formula in the form

$$P(E) = \int_E f(x) dx ,$$

where $f(x)$ is the constant function with value 1.

Density Functions of Continuous Random Variables

Let X be a continuous real-valued random variable. A *density function* for X is a real-valued function f which satisfies

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

for all $a, b \in \mathbf{R}$.

- It is *not* the case that all continuous real-valued random variables possess density functions.
- In terms of the density $f(x)$, if E is a subset of \mathbb{R} , then

$$P(X \in E) = \int_E f(x) dx .$$

Example

- In the spinner experiment, we choose for our set of outcomes the interval $0 \leq x < 1$, and for our density function

$$f(x) = \begin{cases} 1, & \text{if } 0 \leq x < 1, \\ 0, & \text{otherwise.} \end{cases}$$

- If E is the event that the head of the spinner falls in the upper half of the circle, then $E = \{x : 0 \leq x \leq 1/2\}$, and so

$$P(E) = \int_0^{1/2} 1 \, dx = \frac{1}{2}.$$

- More generally, if E is the event that the head falls in the interval $[a, b]$, then

$$P(E) = \int_a^b 1 \, dx = b - a .$$

Example: Continuous Uniform Density

- The simplest density function corresponds to the random variable U whose value represents the outcome of the experiment consisting of choosing a real number at random from the interval $[a, b]$.


$$f(w) = \begin{cases} 1/(b - a), & \text{if } a \leq \omega \leq b \\ 0, & \text{otherwise.} \end{cases}$$

Normal Density

- The *normal density* function with parameters μ and σ is defined as follows:

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2} .$$

- The parameter μ represents the “center” of the density.
- The parameter σ is a measure of the “spread” of the density, and thus it is assumed to be positive.





Central Limit Theorem for Bernoulli Trials

- We deal only with the case that $\mu = 0$ and $\sigma = 1$.
- We will call this particular normal density function the *standard normal density*, and we will denote it by $\phi(x)$:

$$\phi(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} .$$

- Consider a Bernoulli trials process with probability p for success on each trial.
- Let $X_i = 1$ or 0 according as the i th outcome is a success or failure, and let $S_n = X_1 + X_2 + \cdots + X_n$.
- Then S_n is the number of successes in n trials.
- We know that S_n has as its distribution the binomial probabilities $b(n, p, j)$.



Standardized Sums

- We can prevent the drifting of these spike graphs by subtracting the expected number of successes np from S_n .
- We obtain the new random variable $S_n - np$.
- Now the maximum values of the distributions will always be near 0.
- To prevent the spreading of these spike graphs, we can normalize $S_n - np$ to have variance 1 by dividing by its standard deviation \sqrt{npq}

Definition

The *standardized sum* of S_n is given by

$$S_n^* = \frac{S_n - np}{\sqrt{npq}} .$$

S_n^* always has expected value 0 and variance 1.

- We plot a spike graph with the spikes placed at the possible values of S_n^* : x_0, x_1, \dots, x_n , where


$$x_j = \frac{j - np}{\sqrt{npq}} .$$

- We make the height of the spike at x_j equal to the distribution value $b(n, p, j)$.

- We plot a spike graph with the spikes placed at the possible values of S_n^* : x_0, x_1, \dots, x_n , where

$$x_j = \frac{j - np}{\sqrt{npq}} .$$

- We make the height of the spike at x_j equal to the distribution value $b(n, p, j)$.



- Let ε be the distance between consecutive spikes.
- to change the spike graph so that the area under this curve has value 1, we need only multiply the heights of the spikes by $1/\varepsilon$.
- We see that


$$\varepsilon = \frac{1}{\sqrt{npq}} .$$

- Let us fix a value x on the x -axis and let n be a fixed positive integer.
- Then the point x_j that is closest to x has a subscript j given by the formula

$$j = \langle np + x\sqrt{npq} \rangle .$$

- Thus the height of the spike above x_j will be

$$\sqrt{npq} b(n, p, j) = \sqrt{npq} b(n, p, \langle np + x_j\sqrt{npq} \rangle) .$$



Central Limit Theorem for Binomial Distributions

Theorem. *For the binomial distribution $b(n, p, j)$ we have*

$$\lim_{n \rightarrow \infty} \sqrt{npq} b(n, p, \langle np + x\sqrt{npq} \rangle) = \phi(x) ,$$

where $\phi(x)$ is the standard normal density.

Approximating Binomial Distributions

- To find an approximation for $b(n, p, j)$, we set

$$j = np + x\sqrt{npq}$$

- Solve for x

$$x = \frac{j - np}{\sqrt{npq}} .$$

$$\begin{aligned} b(n, p, j) &\approx \frac{\phi(x)}{\sqrt{npq}} \\ &= \frac{1}{\sqrt{npq}} \phi\left(\frac{j - np}{\sqrt{npq}}\right) . \end{aligned}$$

Example

- Let us estimate the probability of exactly 55 heads in 100 tosses of a coin.
- For this case $np = 100 \cdot 1/2 = 50$ and $\sqrt{npq} = \sqrt{100 \cdot 1/2 \cdot 1/2} = 5$.
- Thus $x_{55} = (55 - 50)/5 = 1$ and

$$\begin{aligned} P(S_{100} = 55) &\sim \frac{\phi(1)}{5} = \frac{1}{5} \left(\frac{1}{\sqrt{2\pi}} e^{-1/2} \right) \\ &= .0484 . \end{aligned}$$