

MATH 20, WORKSHEET 5

EDGAR COSTA

DUE FRIDAY OCTOBER 13TH

- (1) If X and Y are any two random variables, then the covariance of X and Y is defined by

$$\text{Cov}[X, Y] := E[(X - E[X])(Y - E[Y])].$$

Note that $\text{Cov}(X, X) = V(X)$. Show that, if X and Y are independent, then $\text{Cov}[X, Y] = 0$; and show, by an example, that we can have $\text{Cov}[X, Y] = 0$ and X and Y not independent.

- (2) For a sequence of Bernoulli trials, let X_1 be the number of trials until the first success. For $j \geq 2$, let X_j be the number of trials after the $(j - 1)$ st success until the j th success. It can be shown that X_1, X_2, \dots is an independent trials process.

- (a) What is the common distribution, expected value, and variance for X_j ? You can use the fact that:

$$x + 4x^2 + 9x^3 + 16x^4 + \dots = x \frac{1+x}{(1-x)^3}$$

or equivalently

$$1 + 4x + 9x^2 + 16x^3 + \dots = \frac{1+x}{(1-x)^3}.$$

- (b) Let $T_n = X_1 + X_2 + \dots + X_n$. Then T_n is the time until the n th success. Find $E[T_n]$ and $V[T_n]$.
- (c) Find the distribution function of T_n . (Challenge: show that adds up to one.)
- (3) The Norwich Beer Company runs a fleet of trucks along the 100 mile road from Hangtown to Dry Gulch. The trucks are old, and are apt to break down at any point along the road with equal probability.
- (a) Where should the company locate a garage so as to minimize the expected distance from a typical breakdown to the garage? In other words, if X is a random variable giving the location of the breakdown, measured, say, from Hangtown, and b gives the location of the garage, what choice of b minimizes $E[|X - b|]$?
- (b) Now suppose X is not distributed uniformly over $[0, 100]$, but instead has density function $f_X(x) = 2x/10000$. Then what choice of b minimizes $E(|X - b|)$?