| NAME: | | |
|-------|--|--|
| | | |

Math 20

Midterm 1 July 14, 2017

Prof. Pantone

INSTRUCTIONS: This is a closed book exam and no notes are allowed. You are not to provide or receive help from any outside source during the exam except that you may ask the instructor for clarification of a problem. You have 120 minutes and you should attempt all problems.

- Print your name in the space provided.
- Calculators or other computing devices are not allowed.
- Except when indicated, you must show all work and give justification for your answer. A correct answer with incorrect work will be considered wrong.

All work on this exam should be completed in accordance with the Dartmouth Academic Honor Principle.

TIPS:

- You don't have numerically expand all answers. For example, you can leave an answer in the form $10! \cdot {5 \choose 3}^2$, rather than 362880000.
- Use scratch paper to figure out your answers and proofs before writing them on your exam.
- Work cleanly and neatly; this makes it easier to give partial credit.

| Problem | Points | Score |
|---------|--------|-------|
| 1 | 24 | |
| 2 | 20 | |
| 3 | 10 | |
| 4 | 16 | |
| 5 | 10 | |
| 6 | 10 | |
| 7 | 10 | |
| 8 | 0 | |
| Total | 100 | |

Section 1: True/False.

- 1.~(24) Choose the correct answer. No justification is required for your answers. No partial credit will be awarded.
- (a) If A and B are mutually exclusive events, then $P(A \cup B) = P(A) + P(B) P(A \cap B)$.

True False

(b) If A and B are independent events, then $P(A \cup B) = P(A) + P(B) - P(A \cap B)$.

True False

(c) If A and B are mutually exclusive events, then $P(A \cap B) = P(A)P(B)$.

True False

(d) If A and B are independent events, then $P(A \cap B) = P(A)P(B)$.

True False

| (e) | The number of ways to line up n people in a row is n^n . | |
|-----|--|--------------------------------|
| | True | False |
| (f) | For any sets A and B , it must be true that $A \subseteq (A \setminus B)$. | |
| | True | False |
| (g) | For any sets A and B , it must be true that $(A \setminus B) \subseteq \overline{B}$ | |
| | True | False |
| (h) | The expected value of a random variable is the numerical occur. | outcome that is most likely to |
| | True | False |

Section 2: Fill in the blank.

- 2. (20) No justification is required for your answers. No partial credit will be awarded.
- (a) There are 100 United States senators, two from each state. If a random group of 50 senators is selected, what is the probability that there will be exactly one from each state?

(b) Suppose that you roll four fair six-sided dice simultaneously. What is the probability that together they form a straight: either $\{1,2,3,4\}$, $\{2,3,4,5\}$, or $\{3,4,5,6\}$?

(c) Let A and B be independent events such that P(A) = 0.2 and P(B) = 0.3. What is $P(A \cup B)$?

(d) A couple decides to have children until they either have three children of the same gender, or until they have four children total, whichever comes first. Find the expected number of children they will have.

Section 3: Free Response.

You must show all work to receive credit.

If you need more space you may use the back of the page. You must clearly indicate on the front of the page that there is more work on the back of the page. Please work neatly.

3. (10) If Juliet gets at least 8 hours of sleep the night before class, then she manages to stay awake for the whole lecture 80% of the time. If Juliet gets between 6 and 8 hours of sleep the night before, she stays awake for the whole lecture 60% of the time. If Juliet gets less than 6 hours of sleep, she only stays awake for the whole lecture 10% of the time.

On any given night, there is a 20% probability that Juliet gets at least 8 hours of sleep, a 60% probability that she gets between 6 and 8 hours of sleep, and a 20% probability that she gets less than 6 hours of sleep.

If Juliet falls asleep in class today, what is the probability that she got at least $\underline{\mathbf{6}}$ hours of sleep the night before?

4. (16) A typical deck of cards contains 52 cards, with 13 cards of each suit (the four suits are \heartsuit , \diamondsuit , \spadesuit , \clubsuit). This question asks about the probability of drawing different poker hands when picking 5 cards randomly from a shuffled deck.

The *order* of the cards of a suit is: Ace, 2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K.

If you have questions about what constitutes a deck of cards or about what the different poker hands mean, please ask me.

(a) A flush is a hand consisting of five cards that all have the same suit. For example, $\{3\spadesuit, 7\spadesuit, 8\spadesuit, J\spadesuit, Q\spadesuit\}$ is a flush. Let P_{flush} be the probability that a randomly drawn five-card poker hand is a flush. Find P_{flush} .

(b) A *straight* is a hand consisting of five cards of consecutive value (see the "order" above). The Ace is allowed to act as either the lowest card (below 2), or the highest card (above K). The suits of the cards are irrelevant. Here are, for example, three different straights:

$$\begin{aligned} & \{A,2,3,4,5\} \\ & \{8,9,10,J,Q\} \\ & \{10,J,Q,K,Ace\} \end{aligned}$$

Let P_{straight} be the probability that a randomly drawn five-card poker hand is a straight. Find P_{straight} .

(c) A poker hand can be both a straight and a flush simultaneously (this is called a straight flush). Find the probability Q that a poker hand is a straight, but not a straight flush.

(d) Let $P_{\rm sad}$ be the probability that your five-card hand has absolutely no poker value. This means: it's not a straight, not a flush, and no two of your cards have the same value (i.e., you have no pair, so all five values of the cards are different). Calculate $P_{\rm sad}$.

5. (10) Let Ω be a finite sample space and let A, B, and C be events. Prove that $P(A)P(B|A)P(C|A\cap B)=P(A\cap B\cap C).$

(Do not use tree diagrams in your proof.)


6. (10) Karen is a pretty good tennis player, and her friend, Henry, offers her \$100 if she can complete the following challenge. To win the money, she has to win two tennis matches in a row out of three total. (Just to clarify, this means she gets the money if she wins Matches 1 and 2, or if she wins Matches 2 and 3, but she does not get the money if she only wins Matches 1 and 3.)

In each of her games, she has to play against either her friend Henry (H) or against the club champion Carlton (C). The options are to play Henry in the first and third games and Carlton in the middle (H,C,H) or to play Carlton in the first and third games and Henry in the middle (C,H,C).

Assume that Carlton is a better tennis player than Henry, and that the outcome of each match is independent of the outcomes of the previous matches. Which of the two match orders—CHC or HCH—gives Karen a better chance at winning the \$100?

(*Hint:* Start by setting p to be the probability that Karen beats Carlton in any given match and q to be the probability that Karen beats Henry in any given match. We're assuming p < q.)

7. (10) Buzz has three wooden boxes, and each box has two drawers, one on each side. Buzz takes three gold rings and three silver rings and distributes one into each drawer: one box gets gold rings in both drawers, one gets silver rings in both drawers, and one gets one gold and one silver. See the picture below.



Now, Woody comes in the room and randomly picks one of the three boxes, and randomly opens one of the two drawers in the box. If that drawer contains a gold ring, then what is the probability that the other drawer also contains a gold ring?

| 8. (0) Bonus: (5 points) Shawn and Gus each flip a fair coin n times. What is the probat that they both flip heads the same number of times? | bility |
|---|--------|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |