

Sample L^AT_EX File

David P. Little

January 7, 2004

Abstract

This document represents the output from the file “sample.tex” once compiled using your favorite L^AT_EX compiler. This file should serve as a good example of the basic structure of a “.tex” file as well as many of the most basic commands needed for typesetting documents involving mathematical symbols and expressions. For more of a description on how each command works, please consult the links found on our course webpage.

1 Lists

1. **First Point (Bold Face)**
2. *Second Point (Italic)*
3. **Third Point (Large Font)**
 - (a) First Subpoint (Small Font)
 - (b) Second Subpoint (Tiny Font)

(c) **Third Subpoint (Huge Font)**

- **Bullet Point (Sans Serif)**
- **CIRCLE POINT (SMALL CAPS)**

2 Equations

2.1 Binomial Theorem

Theorem 1 (Binomial Theorem) *For any nonnegative integer n , we have*

$$(1 + x)^n = \sum_{i=0}^n \binom{n}{i} x^i$$

2.2 Taylor Series

The Taylor series expansion for the function e^x is given by

$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \cdots = \sum_{n \geq 0} \frac{x^n}{n!} \tag{1}$$

2.3 Sets

Theorem 2 For any sets A , B and C , we have

$$(A \cup B) - (C - A) = A \cup (B - C)$$

Proof:

$$\begin{aligned}(A \cup B) - (C - A) &= (A \cup B) \cap (C - A)^c \\ &= (A \cup B) \cap (C \cap A^c)^c \\ &= (A \cup B) \cap (C^c \cup A) \\ &= A \cup (B \cap C^c) \\ &= A \cup (B - C)\end{aligned}$$

□

3 Tables

left justified	center	right justified
1	3.14159	5
2.4678	3	1234
3.4678	6.14159	1239

4 A Picture

