

## Definition

If  $\gamma$  is a smooth curve with admissible parameterization  $z : [a, b] \rightarrow \mathbb{C}$  given by  $z(t) = x(t) + iy(t)$ , then the **length of  $\gamma$**  is

$$\ell(\gamma) := \int_a^b |z'(t)| dt = \int_a^b \sqrt{x'(t)^2 + y'(t)^2} dt$$

If  $\Gamma = \gamma_1 + \cdots + \gamma_n$  is a contour, then

$$\ell(\Gamma) = \ell(\gamma_1) + \cdots + \ell(\gamma_n).$$

## Remark

We know from multivariable calculus that  $\ell(\gamma)$ , and hence  $\ell(\Gamma)$ , is independent of admissible parameterization.

# Ordinary Integrals

## Definition

If  $z(t) = u(t) + iv(t)$  and  $z : [a, b] \rightarrow \mathbb{C}$  is continuous, then we define

$$\int_a^b z(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

Since these are just ordinary integrals, if  $F'(t) = z(t)$ , then

$$\int_a^b z(t) dt = F(t) \Big|_a^b = F(b) - F(a).$$

## Example

If  $z(t) = e^{at}$ , then  $z'(t) = ae^{at}$  for any  $a \in \mathbb{C}$ . Hence

$$\int_0^{\frac{\pi}{2}} e^{2it} dt = \frac{e^{2it}}{2i} \Big|_0^{\frac{\pi}{2}} = i.$$