



# Recent developments on chromatic quasisymmetric functions

Michelle Wachs  
University of Miami

# Chromatic symmetric functions





Let  $C(G)$  be set of proper colorings of graph  $G = ([n], E)$ , where a proper coloring is a map  $c : [n] \rightarrow \mathbb{P}$  such that  $c(i) \neq c(j)$  if  $\{i, j\} \in E$ .

Chromatic symmetric function (Stanley, 1995)

$$X_G(\mathbf{x}) := \sum_{c \in C(G)} x_{c(1)} x_{c(2)} \cdots x_{c(n)}$$

# Chromatic symmetric functions



Let  $C(G)$  be set of proper colorings of graph  $G = ([n], E)$ , where a proper coloring is a map  $c : [n] \rightarrow \mathbb{P}$  such that  $c(i) \neq c(j)$  if  $\{i, j\} \in E$ .

Chromatic symmetric function (Stanley, 1995)

$$X_G(\mathbf{x}) := \sum_{c \in C(G)} x_{c(1)} x_{c(2)} \cdots x_{c(n)}$$

$$X_G(\underbrace{1, 1, \dots, 1}_m, 0, 0, \dots) = \chi_G(m)$$

# Chromatic symmetric functions

Let  $\Pi_G$  be the bond lattice of  $G$ .

Whitney (1932):

$$\chi_G(m) = \sum_{\pi \in \Pi_G} \mu(\hat{0}, \pi) m^{|\pi|}$$

# Chromatic symmetric functions

Let  $\Pi_G$  be the bond lattice of  $G$ .

Whitney (1932):

$$\chi_G(m) = \sum_{\pi \in \Pi_G} \mu(\hat{0}, \pi) m^{|\pi|}$$

Stanley (1995): Let  $p_\lambda$  denote the power-sum symmetric function associated with  $\lambda \vdash n$ . Then

$$X_G(\mathbf{x}) = \sum_{\pi \in \Pi_G} \mu(\hat{0}, \pi) p_{\text{type}(\pi)}(\mathbf{x}).$$

Equivalently

$$\omega X_G(\mathbf{x}) = \sum_{\pi \in \Pi_G} |\mu(\hat{0}, \pi)| p_{\text{type}(\pi)}(\mathbf{x}),$$

which implies that  $\omega X_G(\mathbf{x})$  is  $p$ -positive.

## b-Positivity

Important **bases** for the vector space  $\Lambda_n$  of homogeneous symmetric functions of degree  $n$ :

- complete homogeneous symmetric functions:  $\{h_\lambda : \lambda \vdash n\}$
- elementary symmetric functions:  $\{e_\lambda : \lambda \vdash n\}$
- power-sum symmetric functions:  $\{p_\lambda : \lambda \vdash n\}$
- Schur functions:  $\{s_\lambda : \lambda \vdash n\}$

Involution  $\omega : \Lambda_n \rightarrow \Lambda_n$  defined by  $\omega(h_\lambda) = e_\lambda$ .

Let  $b = \{b_\lambda : \lambda \vdash n\}$  be a basis for  $\Lambda_n$ . A symmetric function  $f \in \Lambda_n$  is said to be **b-positive** if  $f = \sum_{\lambda \vdash n} c_\lambda b_\lambda$ , where  $c_\lambda \geq 0$ .

h-positive  $\implies$  p-positive and Schur-positive.

$f$  is e-positive  $\iff \omega f$  is h-positive.

$$X_{K_{3,1}} = 4e_4 + 5e_{3,1} - 2e_{2,2} + e_{2,1,1}$$

$$X_{K_{3,1}} = 4e_4 + 5e_{3,1} - 2e_{2,2} + e_{2,1,1}$$

- The **incomparability graph**  $\text{inc}(P)$  of a finite poset  $P$  on  $[n]$  is the graph whose edges are pairs of incomparable elements of  $P$ .
- A poset  $P$  is said to be  **$(a + b)$ -free** if  $P$  contains no induced subposet isomorphic to the disjoint union of an  $a$  element chain and a  $b$  element chain.



$$X_{K_{3,1}} = 4e_4 + 5e_{3,1} - 2e_{2,2} + e_{2,1,1}$$

- The **incomparability graph**  $\text{inc}(P)$  of a finite poset  $P$  on  $[n]$  is the graph whose edges are pairs of incomparable elements of  $P$ .
- A poset  $P$  is said to be  **$(a + b)$ -free** if  $P$  contains no induced subposet isomorphic to the disjoint union of an  $a$  element chain and a  $b$  element chain.

Conjecture (Stanley-Stembridge (1993))

*If  $P$  is  $(3 + 1)$ -free then  $X_{\text{inc}(P)}$  is e-positive.*



# Stanley-Stembridge $e$ -positivity conjecture

## Conjecture (Stanley-Stembridge (1993))

*If  $P$  is  $(3+1)$ -free then  $X_{\text{inc}(P)}$  is  $e$ -positive.*

- Gasharov (1994): expansion in the Schur basis  $\{s_\lambda\}$
- Chow (1996): expansion in the fundamental quasisymmetric function basis  $\{F_\mu\}$
- Guay-Paquet (2013): If true for **unit interval orders** (posets that are both  $(3+1)$ -free and  $(2+2)$ -free) then true in general i.e. for posets that are  $(3+1)$ -free.

# Quasisymmetric refinement



Chromatic **quasisymmetric** function (Shareshian and MW)

$$X_G(\mathbf{x}, t) := \sum_{c \in C(G)} t^{\text{des}(c)} x_{c(1)} x_{c(2)} \cdots x_{c(n)}$$

where

$$\text{des}(c) := |\{\{i, j\} \in E(G) : i < j \text{ and } c(i) > c(j)\}|.$$

# Quasisymmetric refinement

$$G = \textcircled{1} \text{---} \textcircled{2} \text{---} \textcircled{3}$$

$$X_G(\mathbf{x}, t) = e_3 + (e_3 + e_{2,1})t + e_3t^2$$

$$G = \textcircled{1} \text{---} \textcircled{3} \text{---} \textcircled{2}$$

$$X_G(\mathbf{x}, t) = (e_3 + F_{1,2}) + 2e_3t + (e_3 + F_{2,1})t^2$$

where  $F_\mu(x_1, x_2, \dots) :=$  fundamental quasisymmetric function indexed by composition  $\mu$

# Chromatic quasisymmetric functions that are symmetric

A **natural unit interval order** is a unit interval order with a certain natural canonical labeling.

**Example:** The poset  $P_{n,r}$  on  $[n]$  with order relation given by  $i <_P j$  if  $j - i \geq r$ . Let

$$G_{n,r} := \text{inc}(P_{n,r}) = ([n], \{\{i, j\} : 0 < j - i < r\})$$

When  $r = 2$ ,  $G_{n,r}$  is the path

$$1 - 2 - \dots - n$$

and

$$X_{G_{n,r}} = \sum_{w \in W_n} t^{\text{des}(w)} x_w,$$

where  $W_n = \{w \in \mathbb{P}^n : \text{adjacent letters of } w \text{ are distinct}\}$ .

# Chromatic quasisymmetric functions that are symmetric

## Theorem (Shareshian and MW)

If  $G$  is the incomparability graph of a natural unit interval order then the coefficients of powers of  $t$  in  $X_G(\mathbf{x}, t)$  are *symmetric functions* and form a *palindromic sequence*.

$$X_{G_{3,2}} = e_3 + (e_3 + e_{2,1})t + e_3t^2$$

$$X_{G_{4,2}} = e_4 + (e_4 + e_{3,1} + e_{2,2})t + (e_4 + e_{3,1} + e_{2,2})t^2 + e_4t^3$$

# Chromatic quasisymmetric functions that are symmetric

## Theorem (Shareshian and MW)

If  $G$  is the incomparability graph of a natural unit interval order then the coefficients of powers of  $t$  in  $X_G(\mathbf{x}, t)$  are *symmetric functions* and form a *palindromic sequence*.

$$X_{G_{3,2}} = e_3 + (e_3 + e_{2,1})t + e_3t^2$$

$$X_{G_{4,2}} = e_4 + (e_4 + e_{3,1} + e_{2,2})t + (e_4 + e_{3,1} + e_{2,2})t^2 + e_4t^3$$

## Conjecture (Shareshian and MW - refinement of Stan-Stem)

If  $G$  is the incomparability graph of a natural unit interval order then the coefficients of powers of  $t$  in  $X_G(\mathbf{x}, t)$  are *e-positive* and form an *e-unimodal sequence*.

True for

$r = 1, n$  (easy)

$r = 2, n - 1, n - 2$  (work of Shareshian and MW)

$1 < r < n \leq 9$  (computer)

## Our approach - a bridge to Hessenberg varieties

Let  $G$  be a **natural unit interval graph** (i.e., the incomparability graph of a natural unit interval order).

Let  $\mathcal{H}_G$  be the regular semisimple Hessenberg variety associated with  $G$ . Tymoczko uses GKM theory to define a representation of  $\mathfrak{S}_n$  on each cohomology  $H^{2j}(\mathcal{H}_G)$ .

### Conjecture (Shareshian and MW (2012))

Let  $\text{ch}H^{2j}(\mathcal{H}_G)$  be the Frobenius characteristic of Tymoczko's representation of  $\mathfrak{S}_n$  on  $H^{2j}(\mathcal{H}_G)$ . Then

$$\omega X_G(\mathbf{x}, t) = \sum_{j \geq 0} \text{ch}H^{2j}(\mathcal{H}_G)t^j.$$



# Our approach - a bridge to Hessenberg varieties

Let  $G$  be a **natural unit interval graph** (i.e., the incomparability graph of a natural unit interval order).

Let  $\mathcal{H}_G$  be the regular semisimple Hessenberg variety associated with  $G$ . Tymoczko uses GKM theory to define a representation of  $\mathfrak{S}_n$  on each cohomology  $H^{2j}(\mathcal{H}_G)$ .

## Conjecture (Shareshian and MW (2012))

*Let  $\text{ch}H^{2j}(\mathcal{H}_G)$  be the Frobenius characteristic of Tymoczko's representation of  $\mathfrak{S}_n$  on  $H^{2j}(\mathcal{H}_G)$ . Then*

$$\omega X_G(\mathbf{x}, t) = \sum_{j \geq 0} \text{ch}H^{2j}(\mathcal{H}_G)t^j.$$

If this conjecture is true then our refinement of the Stanley-Stembridge e-positivity conjecture is equivalent to

## Conjecture

*Tymoczko's representation of  $\mathfrak{S}_n$  on  $H^{2j}(\mathcal{H}_G)$  is a permutation representation for which each point stabilizer is a Young subgroup.*

# The bridge conjecture is true!

Let  $G$  be a natural unit interval graph.

Theorem (Brosnan and Chow (2015), Guay-Paquet (2016))

Let  $\text{ch}H^{2j}(\mathcal{H}_G)$  be the Frobenius characteristic of Tymoczko's representation of  $\mathfrak{S}_n$  on  $H^{2j}(\mathcal{H}_G)$ . Then

$$\omega X_G(\mathbf{x}, t) = \sum_{j \geq 0} \text{ch}H^{2j}(\mathcal{H}_G)t^j.$$

Combinatorial consequences:

- $X_G(\mathbf{x}, t)$  is Schur-positive and **Schur-unimodal**.
- Generalized  $q$ -Eulerian polynomials are  $q$ -unimodal.

Algebro-geometric consequences:

- Multiplicity of irreducibles in Tymoczko's representation can be obtained from the expansion of  $X_G(\mathbf{x}, t)$  in Schur basis.
- Character of Tymoczko's representation can be obtained from expansion of  $X_G(\mathbf{x}, t)$  in power-sum basis.

## Schur and power-sum expansions

Let  $G = ([n], E)$  be a natural unit interval graph, and let  $P$  be such that  $G = \text{inc}(P)$ .

For  $\sigma \in \mathfrak{S}_n$ , a **G-inversion** of  $\sigma$  is an inversion  $(\sigma(i), \sigma(j))$  of  $\sigma$  such that  $\{\sigma(i), \sigma(j)\} \in E$ . Let  $\text{inv}_G(\sigma)$  be the number of  $G$ -inversions of  $\sigma$ .

Theorem (Shareshian and MW,  $t=1$  Gasharov)

$$X_G(\mathbf{x}, t) = \sum_{\lambda \vdash n} \sum_{T \in \mathcal{T}_{P, \lambda}} t^{\text{inv}_G(w(T))} s_\lambda.$$

Consequently  $X_G(\mathbf{x}, t)$  is Schur-positive.

Theorem (Athanasiadis, conjectured by Shareshian and MW)

$$\omega X_G(\mathbf{x}, t) = \sum_{\lambda \vdash n} \sum_{T \in \mathcal{R}_{P, \lambda}} t^{\text{inv}_G(w(T))} z_\lambda^{-1} p_\lambda$$

Consequently  $\omega X_G(\mathbf{x}, t)$  is  $p$ -positive.

A weakly increasing sequence  $\mathbf{m} = (m_1, \dots, m_n)$  of integers satisfying  $1 \leq i \leq m_i \leq n$ , will be called a **Hessenberg sequence**.

Let  $\mathcal{F}_n$  be the set of all flags of subspaces of  $\mathbb{C}^n$

$$F : F_1 \subset F_2 \subset \dots \subset F_n = \mathbb{C}^n$$

where  $\dim F_i = i$ .

Fix an  $n \times n$  diagonal matrix  $D$  with distinct diagonal entries and let  $\mathbf{m} = (m_1, \dots, m_n)$  be a Hessenberg sequence.

The **type A regular semisimple Hessenberg variety associated with  $\mathbf{m}$**  is

$$\mathcal{H}(\mathbf{m}) := \{F \in \mathcal{F}_n \mid DF_i \subseteq F_{m_i} \ \forall i \in [n]\}.$$

A weakly increasing sequence  $\mathbf{m} = (m_1, \dots, m_n)$  of integers satisfying  $1 \leq i \leq m_i \leq n$ , will be called a **Hessenberg sequence**.

Let  $\mathcal{F}_n$  be the set of all flags of subspaces of  $\mathbb{C}^n$

$$F : F_1 \subset F_2 \subset \dots \subset F_n = \mathbb{C}^n$$

where  $\dim F_i = i$ .

Fix an  $n \times n$  diagonal matrix  $D$  with distinct diagonal entries and let  $\mathbf{m} = (m_1, \dots, m_n)$  be a Hessenberg sequence.

The **type A regular semisimple Hessenberg variety associated with  $\mathbf{m}$**  is

$$\mathcal{H}(\mathbf{m}) := \{F \in \mathcal{F}_n \mid DF_i \subseteq F_{m_i} \ \forall i \in [n]\}.$$

Every natural unit interval graph  $G$  can be associated with a Hessenberg sequence  $\mathbf{m}(G)$ . Let

$$\mathcal{H}_G := \mathcal{H}(\mathbf{m}(G)).$$

# GKM theory and moment graphs

Goresky, Kottwitz, MacPherson (1998): Construction of equivariant cohomology ring of smooth complex projective varieties with a torus action. From this one gets ordinary cohomology ring.

The group  $T$  of nonsingular  $n \times n$  diagonal matrices acts on

$$\mathcal{H}_G := \{F \in \mathcal{F}_n \mid DF_i \subseteq F_{m_i(G)} \quad \forall i \in [n]\}.$$

by left multiplication.

**Moment graph:** graph whose vertices are  $T$ -fixed points and edges are one-dimensional orbits.

Fixed points of the torus action:

$$F_\sigma : \langle e_{\sigma(1)} \rangle \subset \langle e_{\sigma(1)}, e_{\sigma(2)} \rangle \subset \cdots \subset \langle e_{\sigma(1)}, \dots, e_{\sigma(n)} \rangle$$

where  $\sigma$  is a permutation.

So the vertices of the moment graph can be represented by permutations.

# Combinatorial description of the moment graph

For any natural unit interval graph  $G = ([n], E)$ , let  $\Gamma(G)$  be the graph with **vertex set**  $\mathfrak{S}_n$  and **edge set**

$$\{\{\sigma, \sigma(i, j)\} : \sigma \in \mathfrak{S}_n \text{ and } \{i, j\} \in E\}.$$

The edges of  $\Gamma(G)$  are **labeled** as follows: edge  $\{\sigma, (a, b)\sigma\}$  is labeled with  $(a, b)$ , where  $a < b$ .

$\Gamma(G)$  is the moment graph for the Hessenberg variety  $\mathcal{H}_G$ .

# Combinatorial description of the moment graph

For any natural unit interval graph  $G = ([n], E)$ , let  $\Gamma(G)$  be the graph with **vertex set**  $\mathfrak{S}_n$  and **edge set**


$$\{ \{\sigma, \sigma(i, j)\} : \sigma \in \mathfrak{S}_n \text{ and } \{i, j\} \in E \}.$$

The edges of  $\Gamma(G)$  are **labeled** as follows: edge  $\{\sigma, (a, b)\sigma\}$  is labeled with  $(a, b)$ , where  $a < b$ .

$\Gamma(G)$  is the moment graph for the Hessenberg variety  $\mathcal{H}_G$ .

**Example:**  $n = 3$ .

Color coded edge labels: **(1,2)** **(2,3)** **(1,3)**






# The equivariant cohomology ring $H_T^*(\mathcal{H}_G)$

$H_T^*(\mathcal{H}_G)$  is isomorphic to a subring of  $R_n := \prod_{\sigma \in \mathfrak{S}_n} \mathbb{C}[t_1, \dots, t_n]$ .

For  $p \in R_n$ , let  $p_\sigma(t_1, \dots, t_n) \in \mathbb{C}[t_1, \dots, t_n]$  denote the  $\sigma$ -component of  $p$ , where  $\sigma \in \mathfrak{S}_n$ .

$p \in R_n$  satisfies the **edge condition for the moment graph  $\Gamma_G$**  if for all edges  $\{\sigma, \tau\}$  of  $\Gamma(G)$  with label  $(i, j)$ , the polynomial  $p_\sigma(t_1, \dots, t_n) - p_\tau(t_1, \dots, t_n)$  is divisible by  $t_i - t_j$ .

Color coded edge labels: (1,2) (2,3) (1,3)




$H_T^*(\mathcal{H}_G)$  is isomorphic to the subring of  $R_n$  whose elements satisfy the edge condition for  $\Gamma_G$ .

# Tymoczko's representation

$\mathfrak{S}_n$  acts on  $p \in H_T^*(\mathcal{H}_G)$  by

$$(\sigma p)_T(t_1, \dots, t_n) = p_{\sigma^{-1}T}(t_{\sigma(1)}, \dots, t_{\sigma(n)})$$

Color coded edge labels: (1,2) (2,3) (1,3)




# Tymoczko's representation

$\mathfrak{S}_n$  acts on  $p \in H_T^*(\mathcal{H}_G)$  by

$$(\sigma p)_T(t_1, \dots, t_n) = p_{\sigma^{-1}T}(t_{\sigma(1)}, \dots, t_{\sigma(n)})$$

Color coded edge labels: (1,2) (2,3) (1,3)




# Tymoczko's representation

$\mathfrak{S}_n$  acts on  $p \in H_T^*(\mathcal{H}_G)$  by

$$(\sigma p)_T(t_1, \dots, t_n) = p_{\sigma^{-1}T}(t_{\sigma(1)}, \dots, t_{\sigma(n)})$$

Color coded edge labels: (1,2) (2,3) (1,3)




# Tymoczko's representation

$\mathfrak{S}_n$  acts on  $p \in H_T^*(\mathcal{H}_G)$  by

$$(\sigma p)_T(t_1, \dots, t_n) = p_{\sigma^{-1}T}(t_{\sigma(1)}, \dots, t_{\sigma(n)})$$

Color coded edge labels: (1,2) (2,3) (1,3)



$$H^*(\mathcal{H}_G) \cong H_T^*(\mathcal{H}_G) / (\langle t_1, \dots, t_n \rangle H_T^*(\mathcal{H}_G))$$

Since  $\langle t_1, \dots, t_n \rangle H_T^*(\mathcal{H}_G)$  is invariant under the action of  $\mathfrak{S}_n$ , the representation of  $\mathfrak{S}_n$  on  $H_T^*(\mathcal{H}_G)$  induces a representation on the graded ring  $H^*(\mathcal{H}_G)$ .


The proofs of  $\omega X_G(\mathbf{x}, t) = \sum_{j \geq 0} \text{ch} H^{2j}(\mathcal{H}_G) t^j$

- **Brosnan and Chow** reduce the problem of computing Tymaczczo's representation of  $\mathfrak{S}_n$  on regular semisimple Hessenberg varieties to that of computing the Betti numbers of regular (but not nec. semisimple) Hessenberg varieties. To do this they use results from the **theory of local systems and perverse sheaves**. In particular they use the local invariant cycle theorem of Beilinson-Bernstein-Deligne
- **Guay-Paquet** introduces a new **Hopf algebra** on labeled graphs to recursively decompose the regular semisimple Hessenberg varieties.

## Other recently discovered connections with $X_G(\mathbf{x}, t)$

- Hecke algebra characters evaluated at Kazhdan-Lusztig basis elements: [Clearman, Hyatt, Shelton and Skandera \(2015\)](#).  
This is a  $t$ -analog of work of Haiman (1993).
- LLT polynomials and Macdonald polynomials: [Haglund and Wilson \(2016\)](#).


# Chromatic quasymmetric function of the cycle graph



Not a unit interval graph



# Chromatic quasisymmetric function of the cycle graph




Not a unit interval graph

Theorem (Stanley (1995))

$$\sum_{n \geq 2} X_{C_n}(\mathbf{x}) z^n = \frac{\sum_{k \geq 2} k(k-1) e_k z^k}{1 - \sum_{k \geq 2} (k-1) e_k z^k}.$$

Consequently  $X_{C_n}(\mathbf{x})$  is  $e$ -positive.

# Chromatic quasisymmetric function of the cycle graph



Not a unit interval graph

Theorem (Stanley (1995))

$$\sum_{n \geq 2} X_{C_n}(\mathbf{x}) z^n = \frac{\sum_{k \geq 2} k(k-1) e_k z^k}{1 - \sum_{k \geq 2} (k-1) e_k z^k}.$$

Consequently  $X_{C_n}(\mathbf{x})$  is  $e$ -positive.

Theorem (Ellzey and MW)

$$\sum_{n \geq 2} X_{C_n}(\mathbf{x}, t) z^n = \frac{\sum_{k \geq 2} ([2]_t [k]_t + k t^2 [k-3]_t) e_k z^k}{1 - t \sum_{k \geq 2} [k-1]_t e_k z^k}$$

Consequently  $X_{C_n}(\mathbf{x}, t)$  is  $e$ -positive.

$t$ -analog:  $[n]_t := 1 + t + \cdots + t^{n-1}$ .

# New improved version


Chromatic **quasisymmetric** function for labeled graphs

$$X_G(\mathbf{x}, t) := \sum_{c \in \mathcal{C}(G)} t^{\text{des}(c)} x_{c(1)} x_{c(2)} \cdots x_{c(n)}$$

where

$$\text{des}(c) := |\{\{i, j\} \in E(G) : i < j \text{ and } c(i) > c(j)\}|.$$

green < yellow < blue



$$\text{des}(c) = 2$$

# New improved version - digraphs


Chromatic **quasisymmetric** function for labeled graphs

$$X_G(\mathbf{x}, t) := \sum_{c \in \mathcal{C}(G)} t^{\text{des}(c)} x_{c(1)} x_{c(2)} \cdots x_{c(n)}$$

where

$$\text{des}(c) := |\{\{i, j\} \in E(G) : i < j \text{ and } c(i) > c(j)\}|.$$

green < yellow < blue




$$\text{des}(c) = 2$$

For a **digraph**  $\vec{G}$

$$\text{des}(c) : |\{(i, j) \in E(\vec{G}) : c(i) > c(j)\}|$$

labeled graphs  $\equiv$  acyclic digraphs


# The directed cycle



Theorem (Ellzey and MW (2017))

$$\sum_{n \geq 2} X_{C_n}(\mathbf{x}, t) z^n = \frac{\sum_{k \geq 2} ([2]_t [k]_t + k t^2 [k-3]_t) e_k z^k}{1 - t \sum_{k \geq 2} [k-1]_t e_k z^k}$$

Consequently  $X_{C_n}(\mathbf{x}, t)$  is  $e$ -positive.




Theorem (Ellzey (2016))

$$\sum_{n \geq 2} X_{\vec{C}_n}(\mathbf{x}, t) z^n = \frac{t \sum_{k \geq 2} k [k-1]_t e_k z^k}{1 - t \sum_{k \geq 2} [k-1]_t e_k z^k}$$

Consequently  $X_{\vec{C}_n}(\mathbf{x}, t)$  is  $e$ -positive.


# The directed cycle



Theorem (Ellzey and MW (2017))

$$\sum_{n \geq 2} X_{C_n}(\mathbf{x}, t) z^n = \frac{\sum_{k \geq 2} ([2]_t [k]_t + k t^2 [k-3]_t) e_k z^k}{1 - t \sum_{k \geq 2} [k-1]_t e_k z^k}$$

Consequently  $X_{C_n}(\mathbf{x}, t)$  is  $e$ -positive.



Theorem (Ellzey (2016))

$$\sum_{n \geq 2} X_{\vec{C}_n}(\mathbf{x}, t) z^n = \frac{t \sum_{k \geq 2} k [k-1]_t e_k z^k}{1 - t \sum_{k \geq 2} [k-1]_t e_k z^k}$$

Consequently  $X_{\vec{C}_n}(\mathbf{x}, t)$  is  $e$ -positive.

The second theorem was used to prove a result on Smirnov words, which implies both theorems.

Theorem (Athanasiadis, conjectured by Shareshian and MW)

Let  $G$  be the incomparability graph of a natural unit interval order  $P$ . Then

$$\omega X_G(\mathbf{x}, t) = \sum_{\lambda \vdash n} \sum_{T \in \mathcal{R}_{P, \lambda}} t^{\text{inv}_G(w(T))} z_\lambda^{-1} p_\lambda$$

Consequently  $\omega X_G(\mathbf{x}, t)$  is  $p$ -positive.


Theorem (Ellzey (2016))

Let  $\vec{G}$  be a digraph for which  $X_{\vec{G}}(\mathbf{x}, t)$  is symmetric. Then

$$\omega X_{\vec{G}}(\mathbf{x}, t) = \sum_{\lambda \vdash n} \sum_{T \in \mathcal{R}_{\vec{G}, \lambda}} t^{\text{inv}_G(w(T))} z_\lambda^{-1} p_\lambda$$



Consequently  $\omega X_{\vec{G}}(\mathbf{x}, t)$  is  $p$ -positive.

# Cyclic version of unit interval digraph







# Cyclic version of unit interval digraph





# Cyclic version of unit interval digraph



circular indifference  
digraph

# Cyclic version of unit interval digraph



circular indifference digraph

## Theorem (Ellzey (2016))

If  $\vec{G}$  is a circular indifference digraph then  $X_{\vec{G}}(\mathbf{x}, t)$  is symmetric.

## Conjecture

If  $\vec{G}$  is a circular indifference digraph then  $X_{\vec{G}}(\mathbf{x}, t)$  is e-positive and e-unimodal.

## Other work on the digraph version of chromatic quasisymmetric functions

- [Awan and Bernardi \(2016\)](#): Tuttle quasisymmetric functions for directed graphs.
- [Alexandersson and Panova \(2016\)](#): connection to LLT polynomials.